arXiv:math/9911080v3 [math.CV] 27 Apr 2001

SOLVABLE MATRIX REPRESENTATIONS OF KÄHLER GROUPS

Alexander Brudnyi*
Department of Mathematics and Statistics
University of Calgary
Calgary, Canada

Abstract

In the paper we study solvable matrix representations of fundamental groups of compact Kähler manifolds (*Kähler groups*). One of our main results is a factorization theorem for such representations. As an application we prove that the universal covering of a compact Kähler manifold with a residually solvable fundamental group is holomorphically convex.

1. Introduction.

1.1. Let M be a compact Kähler manifold. The fundamental group $\pi_1(M)$ of M can be studied via its representations or equivalently in terms of locally constant sheaves on M. The set $Hom(\pi_1(M), GL_n(\mathbb{C}))$ of all homomorphisms of $\pi_1(M)$ into the matrix Lie group $GL_n(\mathbb{C})$ has the natural structure of a complex affine algebraic variety. The group $GL_n(\mathbb{C})$ acts on $Hom(\pi_1(M), GL_n(\mathbb{C}))$ by pointwise conjugation: $(gf)(s) = gf(s)g^{-1}$, $s \in \pi_1(M)$. The study of geometric properties of $Hom(\pi_1(M), GL_n(\mathbb{C}))$ is of interest because of the relation to the problem of classification of Kähler groups and description of complex manifolds which occur as nonramified coverings of a projective manifold. For any $\rho \in Hom(\pi_1(M), GL_n(\mathbb{C}))$ let us denote by X_ρ the union of all the irreducible components containing ρ. Let $D_n(\mathbb{C}) \subset GL_n(\mathbb{C})$ be the subgroup of diagonal matrices. The primary goal of this paper is to prove a factorization theorem for the elements of X_ρ with $\rho \in Hom(\pi_1(M), D_n(\mathbb{C}))$. Before formulating the result let us first introduce some notation.

Let N be a compact Kähler manifold. Hereafter $\alpha_N : N \longrightarrow Alb(N)$ denotes the Albanese map of N and N_a is the normalization of the image $\alpha_N(N)$. We also denote by $\alpha_N^n : N \longrightarrow N_a$ the holomorphic map which covers α_N.

---

*1991 *Mathematics Subject Classification.* Primary 20F34. Secondary 58A10.
Key words and phrases. Kähler group, fundamental group, holomorphic convexity, flat vector bundle, d-gauge transform, cohomology groups, solvable Lie groups.

Theorem 1.1 *Let M be a compact Kähler manifold. There is a finite Galois covering $r : M_H \longrightarrow M$ with an abelian Galois group $H := H(M)$ such that for any $\xi \in X_\rho$ with $\rho \in Hom(\pi_1(M), D_n(\mathbb{C}))$ we have*

$$Ker(\alpha^n_{M_{H*}}) \subset Ker(\xi) \ .$$

Remark 1.2 If $\rho \in Hom(\pi_1(M), D_n(\mathbb{C}))$ is the trivial homomorphism, then our method of proof gives also $Ker(\alpha^n_{M*}) \subset Ker(\xi)$ for any $\xi \in X_\rho$.

Example 1.3 Let $T_n(\mathbb{C}) \subset GL_n(\mathbb{C})$ be the subgroup of complex upper triangular matrices. For any $\rho \in Hom(\pi_1(M), T_n(\mathbb{C}))$ let $d(\rho) \in Hom(\pi_1(M), D_n(\mathbb{C}))$ be the representation defined by the diagonal of ρ. For the holomorphic diagonal matrix $A(z) := diag[1, z, z^2, ..., z^{n-1}]$, $z \in \mathbb{C}$, we set $\rho(z) := A(z)^{-1}\rho A(z)$. Then $\rho(z)$ determines a holomorphic deformation of ρ to $\rho(0) = d(\rho)$. In particular, $\rho \in X_{d(\rho)}$.

We now generalize the above example. Let V_ρ be a flat vector bundle on M constructed by $\rho \in Hom(\pi_1(M), GL_n(\mathbb{C}))$. Clearly, V_ρ can be also considered as a holomorphic vector bundle. Assume that

(a) As a holomorphic bundle, V_ρ is isomorphic to a bundle $V^{\mathcal{O}}_\rho$ with structure group $T_n(\mathbb{C})$.

Let $Gr^*V^{\mathcal{O}}_\rho$ be the associated graded vector bundle with cocycle defined as the diagonal of the cocycle of $V^{\mathcal{O}}_\rho$. Then $Gr^*V^{\mathcal{O}}_\rho$ is the direct sum of rank-1 holomorphic vector bundles on M. Assume that

(b) The first Chern classes of components of the decomposition of $Gr^*V^{\mathcal{O}}_\rho$ belong to $Tor(H^2(M), \mathbb{Z})$.

Denote the class of representations ρ satisfying (a) and (b) by $S^{\mathcal{O}}_n(M)$. Clearly, $Hom(\pi_1(M), T_n(\mathbb{C})) \subset S^{\mathcal{O}}_n(M)$. Let $D^u_n \subset D_n(\mathbb{C})$ be the subgroup of unitary diagonal matrices and $p_t : M_t \longrightarrow M$ be the Galois covering of M with the Galois group $Tor(\pi_1(M)/D\pi_1(M))$. Here $DG = D^1G$ is the derived subgroup of a group G. (We also set $D^iG = DD^{i-1}G$.) Under the assumptions of Theorem 1.1 we prove

Theorem 1.4 *For any $\xi \in S^{\mathcal{O}}_n(M)$ there is $\rho \in Hom(\pi_1(M), D^u_n)$ such that $p^*_t(\xi) \in X_{p^*_t(\rho)}$. Moreover, $r^*(\xi)$ factors through α_{M_H}, that is,*

$$\xi \circ r_* = \xi' \circ \alpha^n_{M_{H*}}$$

for some $\xi' \in Hom(\pi_1(M_{Ha}), GL_n(\mathbb{C}))$.

Remark 1.5 Assume that $\xi \in S^{\mathcal{O}}_n(M)$ is such that $Gr^*V^{\mathcal{O}}_\xi$ is the direct sum of C^∞-trivial rank-1 holomorphic vector bundles (equivalently, all first Chern classes in (b) above equal 0). Then we will show that $\xi \in X_\rho$ for some $\rho \in Hom(\pi_1(M), D^u_n)$.

Let $E_n(M)$ be the class of homomorphisms $\rho : \pi_1(M) \longrightarrow D_n(\mathbb{C})$ whose diagonal characters ρ_{ii} satisfy $\rho_{ii} = \exp(\widetilde{\rho}_{ii})$ for some $\widetilde{\rho}_{ii} \in Hom(\pi_1(M), \mathbb{C})$. By $S_n(M)$ we denote the class of representations $\rho \in Hom(\pi_1(M), T_n(\mathbb{C}))$ such that the homomorphism $d(\rho)$ defined by the diagonal of ρ belongs to $E_n(M)$. If $\pi : \widetilde{T_n(\mathbb{C})} \longrightarrow T_n(\mathbb{C})$ is the universal covering, then, according to [O, Th.5.5], $\rho \in S_n(M)$ if and only if $\rho = \pi \circ \widetilde{\rho}$ for some $\widetilde{\rho} \in Hom(\pi_1(M), \widetilde{T_n(\mathbb{C})})$. In Lemma 4.6 we will show that $X_\rho \subset X_{d(\rho)}$ for any $\rho \in S_n(M)$. Let $T_n^u \subset T_n(\mathbb{C})$ be the subgroup of matrices with unitary diagonals and $\widetilde{T_n^u} \subset \widetilde{T_n(\mathbb{C})}$ be the universal covering of T_n^u. We set $S_n^u(M) := S_n(M) \cap Hom(\pi_1(M), T_n^u)$ and define

$$G_s^0(M) = \bigcap_{n\geq 1} \bigcap_{\rho\in S_n(M)} Ker(\rho) \qquad G_{su}^0(M) = \bigcap_{n\geq 1} \bigcap_{\rho\in S_n^u(M)} Ker(\rho).$$

Notice that the above groups are kernels of $\mathcal{P}$-completion homomorphisms $\eta^{\mathcal{P}} : \pi_1(M) \longrightarrow \pi_1(M)^{\mathcal{P}}$ where $\mathcal{P}$ is the category of groups $\widetilde{T_n(\mathbb{C})}$ and $\widetilde{T_n^u}$, respectively. In general we can say only that $G_s^0(M) \subseteq G_{su}^0(M)$. However, if M is a compact Kähler manifold then

Theorem 1.6 $G_s^0(M) = G_{su}^0(M)$.

1.2. We apply Theorem 1.1 to the problem of the holomorphic convexity of a Galois covering of a compact Kähler manifold. Central to the subject is a conjecture of Shafarevich according to which the universal covering M_u of a projective manifold M is holomorphically convex, meaning that for every infinite sequence of points without limit points in M_u there exists a holomorphic function unbounded on this sequence. In recent years many new powerful methods have been developed to investigate the structure of the fundamental groups and universal coverings of complex projective manifolds. These methods lead to many new remarkable results. In particular quite a few positive results on the Shafarevich's conjecture were proved, e.g. the results of F. Campana, H. Grauert, R. Gurjar, L. Katzarkov, M. Nori, M. Ramachandran, C. Simpson, S. Shasrty, K. Zuo, S.T. Yau (see e.g. [C], [Ka1,2], [KaR], [Ko1,2], [N], [S], [Y]). It was also shown by F.Bogomolov and L.Katzarkov [BK] that a counterexample to the Shafarevich conjecture exists provided that one can answer affirmatively a certain group theoretic question.

For a compact Kähler manifold M we define

$$G_s(M) := \bigcap_{n\geq 1} \bigcap_{\rho\in Hom(\pi_1(M),T_n(\mathbb{C}))} Ker(\rho) \tag{1.1}$$

and let M_s be the regular covering of M with the transformation group $\pi_1(M)/G_s(M)$.

Theorem 1.7 *M_s is a holomorphically convex manifold. Thus if $p : M_1 \longrightarrow M$ is a finite unbranched covering of M such that the family $\{Hom(\pi_1(M_1), T_n(\mathbb{C}))\}_{n\geq 1}$ separates the elements of $\pi_1(M_1)$ then the universal covering M_u of M is holomorphically convex.*

As a corollary we obtain the result of Katzarkov [Ka1] on the holomorphic convexity of the Malcev covering of any smooth projective manifold.

Corollary 1.8 *Let M be a compact Kähler manifold such that $G_s(M) = \{e\}$ and $\pi_2(M) = 0$. Then M_u is a Stein manifold.*

1.3. We formulate several corollaries on the structure of Kähler groups.

Let C be a smooth compact complex curve of genus $g \geq 1$. Any character $\xi \in Hom(\pi_1(C), \mathbb{C}^*)$ can be written as $\exp(\widetilde{\xi})$ for some $\widetilde{\xi} \in Hom(\pi_1(C), \mathbb{C})$. In turn, $\widetilde{\xi}$ is defined by integration along the paths of a harmonic 1-form ω on C. Let $\omega = \omega' + \omega''$ be the type decomposition of ω into the sum of holomorphic and antiholomorphic 1-forms and $\widetilde{\xi}'' \in Hom(\pi_1(C), \mathbb{C})$ be the homomorphism defined by integration of $\omega'' - \overline{\omega''}$. We set $\xi''_u = \exp(\widetilde{\xi}'')$. Then $\xi''_u \in Hom(\pi_1(C), U_1)$ where U_1 is the unitary group. Notice also that ξ''_u is uniquely defined by ξ. Let $f : M \longrightarrow C$ be a holomorphic surjective map with connected fibres of a compact Kähler manifold M onto a smooth compact complex curve C of genus $g \geq 1$.

Theorem 1.9 *Assume that $\rho \in Hom(\pi_1(M), T_n(\mathbb{C}))$ satisfies*

$$(1) \quad d(\rho) = \bigoplus_{i=1}^{n} f^*(\rho_i) \quad \text{for some} \quad \rho_i \in Hom(\pi_1(C), \mathbb{C}^*);$$

$$(2) \quad \rho''_{iu} \otimes (\rho''_{ju})^{-1} \quad \text{is not a torsion character for any} \quad i \neq j.$$

Then there is $\widetilde{\rho} \in Hom(\pi_1(C), T_n(\mathbb{C}))$ such that $\rho = f^*(\widetilde{\rho})$.

Remark 1.10 If $\rho \in Hom(\pi_1(M), T^u_n)$ then (2) acquires the form $\rho_i \otimes \rho_j^{-1}$ is not torsion for any $i \neq j$.

Let G be the fundamental group of a compact Riemann surface of genus $g \geq 2$ and G_1 be an extension of G by $\mathbb{Z}^k$, $k \geq 1$, such that the conjugate action s of G on $\mathbb{Z}^k$ is trivial on DG. Thinking of $\mathbb{Z}^k$ as a subgroup of $\mathbb{Z}^k \otimes \mathbb{C}$ we can extend s to a representation $s' : G \longrightarrow GL_k(\mathbb{C})$ so that $s'(g)|_{\mathbb{Z}^k} = s(g)$ for any $g \in G$. Then $DG \subset Ker(s')$ and so s' admits a decomposition $\oplus_{j=1}^m s_j$ where s_j is equivalent to a nilpotent representation $G \longrightarrow T_{k_j}(\mathbb{C})$ with diagonal character ρ_j.

Theorem 1.11 *Assume that a Kähler group F is defined as*

$$\{e\} \longrightarrow K \xrightarrow{i} F \xrightarrow{j} G_1 \longrightarrow \{e\}$$

and $j^{-1}(\mathbb{Z}^k) \subset F$ does not admit a surjective homomorphism onto a free group with countable number of generators. Then all characters ρ_j are torsion.

The above theorem produces many examples of non-Kähler groups even in the case when K is finitely presented. For instance, a semidirect product of G with $\mathbb{Z}^k$ defined by $s : G \longrightarrow SL_k(\mathbb{Z})$ as above such that not all ρ_j are torsion is not a Kähler group.

We will complete this section by

Theorem 1.12 *Let M be a compact Kähler manifold satisfying the conditions of Corollary 1.8 and M_H be the same as in Theorem 1.1. Assume also that*

$$dim_{\mathbb{C}}M \geq \frac{1}{2}rank(\pi_1(M)/D^2\pi_1(M)) \ .$$

Then $dim_{\mathbb{C}}M = \frac{1}{2}rank(\pi_1(M)/D^2\pi_1(M))$ and $\pi_1(M_H)$ is a free abelian group. If, in addition,

$$\pi_i(M_u) = 0 \text{ for } 1 \leq i \leq dim_{\mathbb{C}}M$$

then $\alpha_{M_H} : M_H \longrightarrow Alb(M_H)$ is biholomorphic.

Remark 1.13 If $\pi_1(M)$ is residually torsion free nilpotent satisfying $\pi_2(M) = 0$ and $dim_{\mathbb{C}}M \geq \frac{1}{2}rank(\pi_1(M)/D\pi_1(M))$ then $dim_{\mathbb{C}}M = \frac{1}{2}rank(\pi_1(M)/D\pi_1(M))$ and $\pi_1(M)$ is free abelian. If, in addition, $\pi_i(M_u) = 0$ for $1 \leq i \leq dim_{\mathbb{C}}M$ then α_M is biholomorphic.

In particular, any compact complex manifold M with a nonzero Kobayashi semimetric satisfying the conditions of Theorem 1.12 or Remark 1.13 is not Kähler.

In the next section we reduce Theorem 1.1 to the case of representations from $Hom(\pi_1(M), T_n(\mathbb{C}))$. Also we describe a classification theorem for a class of flat connections on $M \times \mathbb{C}^n$ with triangular (0,1)-components (see [Br]) which is crucial in the proof of Theorem 1.1.

2. Flat Connections and Flat Vector Bundles.

2.1. In this section we collect some results on the relation between flat vector bundles and the flat connections which determine them.

Let M be a compact Kähler manifold and $gl_n(\mathbb{C})$ be the Lie algebra of $GL_n(\mathbb{C})$. Consider the family of C^∞-trivial flat vector bundles on M of complex rank n. It is well-known that every bundle from this family is determined by a flat connection on the trivial bundle $M \times \mathbb{C}^n$, that is, by a $gl_n(\mathbb{C})$-valued differential 1-form ω on M satisfying

$$d\omega - \omega \wedge \omega = 0 \ . \qquad (2.1)$$

Denote the class of flat connections by $\mathcal{A}_n(M)$. The group $C^\infty(M, GL_n(\mathbb{C}))$ acts by d-gauge transforms on the set $\mathcal{A}_n(M)$:

$$d_g(\alpha) = g^{-1}\alpha g - g^{-1}dg, \quad g \in C^\infty(M, GL_n(\mathbb{C})), \ \ \alpha \in \mathcal{A}_n(M) \ . \qquad (2.2)$$

Denote the corresponding quotient set by $\mathcal{B}_n(M)$. (We regard $\mathcal{B}_n(M)$ as the set of d-gauge equivalent classes of connections from $\mathcal{A}_n(M)$.) Let $p : M_u \longrightarrow M$ be the universal covering of M and $p^*(\omega)$ be the pullback of ω. Fix a point $x \in M_u$. Then there is a unique solution $I_x(\omega) \in C^\infty(M_u, GL_n(\mathbb{C}))$ of the equation $dF = p^*(\omega)F$ satisfying $I_x(\omega)(x) = E_n$ (here E_n denotes the unit matrix). $I_x(\omega)$ can be obtained by the Picard method of successive approximations (also called *iterated* *path integration*). Let $|\cdot|$ denote the l_2-norm in the space of $n \times n$ matrices. Then we also have

$$|I_x(\omega)(y)| \le e^{cr(y)|\omega|_M} \tag{2.3}$$

where r is the distance from x in the metric pulled back from M, $c := c(M) > 0$ is a constant and $|\cdot|_M$ is an L_2-norm on the space of $gl_n(\mathbb{C})$-valued differential 1-forms on M. Moreover, if the form $\omega = \omega(z)$ depends holomorphically on a parameter z varying in the unit disk $\mathbb{D} \subset \mathbb{C}$, then $I_x(\omega(z))(y)$ is holomorphic in z for any $y \in M_u$.

Let $o(x) := \{h(x)\}_{h\in\pi_1(M)}$ be the orbit of x under the action of the covering group $\pi_1(M)$. Then the restriction $I_x(\omega)|_{o(x)}$ determines a homomorphism of $\pi_1(M)$ into $GL_n(\mathbb{C})$:

$$I_x(\omega)((h_1h_2)(x)) = I_x(\omega)(h_1(x)) \cdot I_x(\omega)(h_2(x)), \quad h_1, h_2 \in \pi_1(M)\ .$$

If $x, y \in M_u$ are distinct points then there is a matrix $C(x, y) \in GL_n(\mathbb{C})$ such that $I_y(\omega)|_{o(y)} = C(x, y)^{-1} I_x(\omega)|_{o(x)} C(x, y)$. Further, for any $g \in C^\infty(M, GL_n(\mathbb{C}))$ there is a matrix $C \in GL_n(\mathbb{C})$ such that

$$I_x(d_g(\alpha)) = C^{-1} I_x(\alpha) C, \quad \alpha \in \mathcal{A}_n(M)\ .$$

Thus I_x determines an injective map $\mathcal{I}_x : \mathcal{B}_n(M) \to Hom(\pi_1(M), GL_n(\mathbb{C}))/GL_n(\mathbb{C})$. Its image consists of equivalence classes of homomorphisms whose associated flat vector bundles are C^∞-trivial.

Another way to construct a representation by a flat connection $\omega \in \mathcal{A}_n(M)$ is as follows (see also, e.g. [O, Sec.5,6]). Let $(U_i)_{i\in I}$ be an open covering of M by sets diffeomorphic to Euclidean balls and let $f_i \in C^\infty(M, GL_n(\mathbb{C}))$ be a solution of $df = \omega f$ on U_i. If we set $c_{ij} = f_i^{-1} f_j$ on $U_i \cap U_j$, then $\{c_{ij}\}$ is a locally constant cocycle and so it determines a flat vector bundle $E(\omega)$ on M. Then a standard procedure (see, e.g. [KN, Ch.2, Sec.9]) allows to construct a $\rho \in Hom(\pi_1(M), GL_n(\mathbb{C}))$ by $E(\omega)$. Notice that ρ coincides with $I_x(\omega)|_{o(x)}$ for a suitable $x \in M_u$.

Let E be a C^∞-trivial flat vector bundle on M. Consider the flat vector bundle $End(E)$. Let $\{c_{ij}\}$ be a locally constant cocycle on an open acyclic covering $(U_i)_{i\in I}$ of M determining E. A family $\{\eta_i\}_{i\in I}$ of matrix-valued p-forms satisfying

$$\eta_j = c_{ij}^{-1} \eta_i c_{ij} \quad \text{on} \quad U_i \cap U_j \tag{2.4}$$

is, by definition, a p-form with values in the bundle $End(E)$. According to (2.4) the operators d and $\wedge$ are well-defined on the set of matrix-valued p-forms. In particular, one can consider $gl_n(\mathbb{C})$-valued 1-forms α with values in $End(E)$ satisfying the flatness condition $d\alpha - \alpha \wedge \alpha = 0$. Let h be a linear C^∞-automorphism of E determined by the family $\{h_i\}_{i\in I}$, ($h_i \in C^\infty(U_i, GL_n(\mathbb{C}))$), satisfying

$$h_j = c_{ij}^{-1} h_i c_{ij} \quad \text{on} \quad U_i \cap U_j\ .$$

Then a d-gauge transform d_h^E defined on the set of $gl_n(\mathbb{C})$-valued 1-forms α with values in $End(E)$ is given by the formula

$$d_h^E(\alpha) = h^{-1}\alpha h - h^{-1} dh\ .$$

Clearly, d_h^E preserves the class of 1-forms satisfying the flatness condition. Let ψ be a flat connection on $M \times \mathbb{C}^n$ determining E and $g_i \in C^\infty(U_i, GL_n(\mathbb{C}))$ be a solution of $df = \psi f$ such that $g_i^{-1} g_j = c_{ij}$. Consider the map τ_ψ defined on the set of matrix-valued 1-forms α on M by

$$\tau_\psi(\alpha_i) = g_i^{-1}(\alpha_i - \psi_i) g_i, \quad \alpha_i := \alpha|_{U_i}, \quad \psi_i := \psi|_{U_i}\ . \tag{2.5}$$

Then τ_ψ maps the set $\mathcal{A}_n(M)$ isomorphically onto the set of $End(E)$-valued matrix 1-forms satisfying the flatness condition. Further, if $h \in C^\infty(M, GL_n(\mathbb{C}))$ then the family $\{g_i^{-1} h g_i\}_{i\in I}$ determines an element of $Aut(E)$. In what follows we identify $C^\infty(M, GL_n(\mathbb{C}))$ with $Aut(E)$. Then we have

$$\tau_\psi \circ d_h = d_h^E \circ \tau_\psi \quad \text{for every} \quad h \in C^\infty(M, GL_n(\mathbb{C}))$$

(for the proof, see e.g. [Br, Prop.2.2]).

2.2. In this part we describe some results of [Br] (see also [BO] for the general case). Let ω be a flat connection on $M \times \mathbb{C}^n$ such that the $(0,1)$-component ω'' of ω is an upper triangular matrix form. Doing a d-gauge transform with a diagonal matrix-function we may assume without loss of generality that $diag(\omega'')$ is a matrix-valued harmonic $(0,1)$- form on M, in particular, that it is d-closed (M is Kähler). Consider another flat connection on $M \times \mathbb{C}^n$ defined by the system of ODEs

$$df = \psi f \tag{2.6}$$

where $\psi := diag(\omega'') - diag(\overline{\omega''})$. Let $\{f_i\}_{i\in I}$, $f_i \in C^\infty(M, D_n^u)$, be a family of local solutions of (2.6) defined on an open acyclic covering $(U_i)_{i\in I}$ of M, where $D_n^u \subset U_n$ is the Lie subgroup of unitary diagonal $n \times n$ matrices. Then the cocycle

$$c_{ij} := f_i^{-1} f_j \quad \text{on} \quad U_i \cap U_j$$

determines a flat vector bundle V_ψ which is the direct sum of C^∞-trivial complex rank-1 flat vector bundles with unitary structure group. In what follows V_ψ-valued harmonic forms are defined with respect to Laplacians constructed by the natural flat Hermitian metric on $End(V_\psi)$. Consider now the map τ_ψ defined in Section 2.1 with ψ as above. Clearly $\tau_\psi(\omega)$ is an $End(V_\psi)$-valued 1-form with a nilpotent (0,1)-component. Recall that a family $\{\eta_i\}_{i\in I}$ of matrix-valued 1-forms satisfying

$$\eta_j = c_{ij}^{-1} \eta_i c_{ij} \quad \text{on} \quad U_i \cap U_j$$

is a nilpotent 1-form with values in $End(V_\psi)$ if every η_i takes its values in the Lie algebra of the Lie group of upper triangular unipotent matrices. It was proved in [Br, Prop.2.3] that

(A) *There is a $g \in C^\infty(M, T_n(\mathbb{C}))$ such that $\tau_\psi(d_g(\omega)) = \eta_1 + \eta_2$, where the (1,0)-form η_1 is ∂-closed and η_2 is a d-closed nilpotent antiholomorphic 1-form. In particular, if the form ω is upper triangular then η_1 is also upper triangular.*

This implies that η_1 can be decomposed into the sum $\alpha + \partial h$, where α is its harmonic component in the Hodge decomposition. Then the Hodge decomposition and the $\partial\overline{\partial}$-lemma (see e.g. [ABCKT, Lm.7.39]) imply

(B) *α is d-harmonic and $(\alpha+\eta_2)\wedge(\alpha+\eta_2)$ represents 0 in the de Rham cohomology group $H^2(M, End(V_\psi))$.*

Conversely, let α be an $End(V_\psi)$-valued d-harmonic $(1,0)$-form and let θ be a d-harmonic nilpotent $End(V_\psi)$-valued $(0,1)$-form. Then we have (see [Br, Prop.2.4]):

(C) *Let $(\alpha+\theta)\wedge(\alpha+\theta)$ represent 0 in $H^2(M, End(V_\psi))$. Then there exists a unique (up to a flat additive summand) section h such that $\alpha+\partial h+\theta$ satisfies the flatness condition. In particular, there is a flat connection ω on $M\times\mathbb{C}^n$ with an upper triangular $(0,1)$-component such that $\tau_\psi(\omega)=\alpha+\partial h+\theta$. If, in addition, α is upper triangular then ω is also upper triangular.*

The following examples will be used in the sequel.

Example 2.1 1. Let C be a compact complex curve of genus $g\geq 1$. Any representation $\rho\in Hom(\pi_1(C), T_n(\mathbb{C}))$ determines a C^∞-trivial flat vector bundle on C. Let $d(\rho)\in Hom(\pi_1(C), D_n(\mathbb{C}))$ be the representation defined by the diagonal of ρ. Then $d(\rho)$ is defined by the system of ODEs $df=\phi f$ with diagonal harmonic 1-form ϕ. Let $\phi=\phi'+\phi''$ be the type decomposition. Denote by $d(\rho)''_u\in Hom(\pi_1(C), D_n^u)$ the representation defined by equation $df=(\phi''-\overline{\phi''})f$. Let $V_{d(\rho)''_u}$ be the flat vector bundle associated to $d(\rho)''_u$. Then $V_{d(\rho)''_u}=\oplus_{i=1}^n V_i$ where each V_i is a complex rank-1 flat vector bundle on C with structure group U_1. Further, $End(V_{d(\rho)''_u})=\oplus_{i,j=1}^n V_i^*\otimes V_j$. Assume that $V_i^*\otimes V_j$ is not trivial for any $i\neq j$. Then it is to see (e.g. by duality) that the de Rham cohomology group $H^2(C, V_i^*\otimes V_j)$ is zero. According to the above results the equivalence class of ρ is defined by an upper triangular harmonic 1-form η with values in $End(V_{d(\rho)''_u})$ such that $\eta\wedge\eta$ represents 0 in $H^2(C, End(V_{d(\rho)''_u}))$ and the (0,1)-component of η is nilpotent. Notice that the identity $H^2(C, V_i^*\otimes V_j)=0$ for $i\neq j$ implies that the first condition for η is always fulfilled. Conversely, any upper-triangular $End(V_{d(\rho)''_u})$-valued harmonic 1-form η with a nilpotent (0,1)-component determines the equivalence class of a representation from $Hom(\pi_1(C), T_n(\mathbb{C}))$.

2. Let M be a compact Kähler manifold, $\rho\in S_n^{\mathcal{O}}(M)$ and $p_t: M_t\longrightarrow M$ be the Galois covering with the Galois group $Tor(\pi_1(M)/D\pi_1(M)$. Let V_ρ be the associated flat vector bundle. Then according to condition (b) in the definition of $S_n^{\mathcal{O}}(M)$, the first Chern classes of components of the decomposition of $Gr^*(p_t^*V_\rho^{\mathcal{O}})$ are zero. Therefore $p_t^*V_\rho$ is defined by a flat connection on $M_t\times\mathbb{C}^n$ with a triangular (0,1)-component and the results of this section describe the equivalence class of $p_t^*V_\rho$.

2.3. In this section we reduce Theorem 1.1 to the case of solvable matrix representations.

Let $p_1(g_1,...,g_k)=e,...,p_l(g_1,...,g_k)=e$ be a finite presentation of $\pi_1(M)$. Here $g_1,...,g_k\in\pi_1(M)$ is a set of generators. Then $Hom(\pi_1(M), GL_n(\mathbb{C}))$ is defined (up to polynomial isomorphisms) as an affine algebraic subvariety of $(GL_n(\mathbb{C}))^k$ by equations

$$p_1(X_1,...,X_k)=E_n,...,p_l(X_1,...,X_k)=E_n,\qquad X_1,...,X_k\in GL_n(\mathbb{C})\ .$$

Let $\rho\in Hom(\pi_1(M), GL_n(\mathbb{C}))$. Assume that there is an irreducible component $X\subset Hom(\pi_1(M), GL_n(\mathbb{C}))$ of ρ such that $dim_{\mathbb{C}}X\geq 1$. Since X is a complex algebraic variety of pure dimension ≥ 1, resolution of singularities shows that there is a non-constant holomorphic map $f : \mathbb{D} \longrightarrow X$ such that $f(0) = \rho$. Let $\mathcal{F}_\rho$ be the family of all such maps. Then there is an open neighbourhood $O \subset X$ of ρ such that for any $w \in O$ there is an $f \in \mathcal{F}_\rho$ so that $w \in f(\mathbb{D})$. Any $f \in \mathcal{F}_\rho$ determines a holomorphic deformation of ρ (which we denote by the same letter). In particular, we have $f(z)(g) = \sum_{i=0}^{\infty} f_i(g)z^i$, $g \in \pi_1(M)$, such that $f(z) \in Hom(\pi_1(M), GL_n(\mathbb{C}))$, $f_0 = \rho$, f_i are functions on $\pi_1(M)$ with values in $gl_n(\mathbb{C})$ and the series converges uniformly on each compact subset of $\mathbb{D}$ for any $g \in \pi_1(M)$.

Let $\mathbb{C}[[z]]$ be the ring of formal power series over $\mathbb{C}$. Let $I_m \subset \mathbb{C}[[z]]$ denote the ideal generated by z^m. Let $\pi_m : \mathbb{C}[[z]] \longrightarrow \mathbb{C}[[z]]/I_m =: A_m[z]$ be the quotient homomorphism. For any $h \in \mathbb{C}[[z]]$ we identify $\pi_m(h)$ with the first m terms of the expansion of h. Let $GL_n(A_m[z])$ be the group of invertible $n \times n$ matrices with entries from $A_m[z]$. Any $a \in GL_n(A_m[z])$ can be written as $a = \sum_{i=0}^{m-1} a_i z^i$, where $a_i \in gl_n(\mathbb{C})$. The element a is invertible if and only if $a_0 \in GL_n(\mathbb{C})$. Similarly, denote by $gl_n(A_m[z])$ the algebra of all $n \times n$ matrices with entries from $A_m[z]$. The homomorphism π_m induces a similar homomorphism (denoted by the same letter) $GL_n(\mathbb{C})[[z]] \longrightarrow GL_n(A_m[z])$ of the space of formal matrix power series which associate to each series the first m terms of its Taylor expansion. Assume now that $f \in \mathcal{F}_\rho$ is a homomorphism depending holomorphically on $z \in \mathbb{D}$. Then $\pi_m(f) \in Hom(\pi_1(M), GL_n(A_m[z]))$. Obviously we have

Proposition 2.2

$$\bigcap_{z \in \mathbb{D}} Ker(f(z)) = \bigcap_{i \geq 0} Ker(\pi_i(f(z))) \qquad \square$$

The main result of this section is the following reduction in Theorem 1.1. We retain the notation introduced in Theorem 1.1.

Proposition 2.3 *Theorem 1.1 is a consequence of the following result:*
For any $\rho \in Hom(\pi_1(M), T_n(\mathbb{C}))$, $n \geq 1$, *we have* $Ker(\alpha^n_{M_{H^*}}) \subset Ker(\rho)$.

Proof. Let $S_n^d \subset gl_n(\mathbb{C})[[z]]$ be the subalgebra of matrix power series $a = \sum_{i=0}^{\infty} a_i z^i$ with diagonal a_0. We set $F_{m,n} := \pi_m(S_n^d) \subset gl_n(A_m[z])$. Any element $b \in F_{m,n}$ admits an expansion $b = \sum_{i=0}^{m-1} b_i z^i$ with diagonal b_0. Further, $F_{m,n}$ is a finite-dimensional Lie algebra over $\mathbb{C}$ with the bracket: $[a, b] := ab - ba$, $a, b \in F_{m,n}$. Let $F_{m,n}^* := \{b \in F_{m,n} : \ b = \sum_{i=0}^{m-1} b_0 z^i, \ b_0 \in D_n\}$ denote the group of invertible elements of $F_{m,n}$ (with respect to multiplication). Then $F_{m,n}^*$ is a complex solvable Lie group. Let $\phi : F_{m,n}^* \longrightarrow GL_p(\mathbb{C})$, $p = dim_{\mathbb{C}} F_{m,n}$, be the faithful representation obtained by the left multiplication of the elements of $F_{m,n}$ by the elements of $F_{m,n}^*$. We set $V_k := \{v \in F_{m,n} : \ v = \sum_{i=k}^{m-1} v_i z^i\}$. Then

$$F_{m,n} = V_0 \supset V_1 \supset V_2 \supset ... \supset V_m = \{0\}$$

is a chain of invariant $F_{m,n}^*$-submodules with respect to ϕ. The quotient representation $F_{m,n}^* \longrightarrow GL(V_{i+1}/V_i)$ is defined by left multiplication of matrices from $gl_n(\mathbb{C})$ by invertible diagonal matrices. In particular, ϕ is equivalent to a representation $\widetilde{\phi} : F_{m,n}^* \longrightarrow T_p(\mathbb{C})$ (defined by the choice of a suitable basis in $F_{m,n}$).

Consider now $\rho \in Hom(\pi_1(M), D_n(\mathbb{C}))$. Let $f \in \mathcal{F}_\rho$ be a holomorphic deformation of ρ. Then $\pi_m(f) \in Hom(\pi_1(M), F^*_{m,n})$. Further, consider $f_m := \widetilde{\phi} \circ \pi_m(f(z))$. It is easy to see that the diagonal characters of f_m are the same as for ρ. Clearly $f_m \in Hom(\pi_1(M), T_p(\mathbb{C}))$. Assume now that Theorem 1.4 is valid for any f_m, $m \geq 0$, and any $n \geq 1$. Then according to Proposition 2.2, $Ker(f(z))$ contains $Ker(\alpha^n_{M_{H*}})$ for any $z \in \mathbb{D}$.

Let $X_1 \subset X_\rho$ be an irreducible component of $Hom(\pi_1(M), GL_n(\mathbb{C}))$ containing ρ and $O \subset X_1$ be an open neighbourhood of ρ such that for any $w \in O$, $w \neq \rho$, there is $f \in \mathcal{F}_\rho$ satisfying $w \in f(\mathbb{D})$. Then the assumption of the proposition and the previous arguments applied to such $f(z)$ imply that $Ker(\alpha^n_{M_{H*}}) \subset Ker(w)$ for any $w \in O$. Let $g \in Ker(\alpha^n_{M_{H*}})$ and $g = p(g_1, ..., g_k)$ be a word representing g. Here $g_1, ..., g_k \in \pi_1(M)$ are generators. Then the complex algebraic matrix function $p(X_1, ..., X_k)$, $X_1, ..., X_k \in GL_n(\mathbb{C})$, equals 0 on an open set $O \subset X_1$. Since X_1 is irreducible, $p \equiv 0$ on X_1. This argument applied to each $g \in Ker(\alpha^n_{M_{H*}})$ shows that $Ker(\alpha^n_{M_{H*}}) \subset Ker(q)$ for any $q \in X_1$. The same is valid for other irreducible components of X_ρ and for any $\rho \in Hom(\pi_1(M), D_n(\mathbb{C}))$. This implies the required statement. $\square$

3. Fibering Kähler Manifolds.

In this section we study some properties of fundamental groups of compact Kähler manifolds which admit surjective holomorphic maps with connected fibers onto compact Riemann surfaces. We also prove Theorems 1.9 and 1.11.

3.1. Let $f : M \longrightarrow C$ be a holomorphic surjective map with connected fibres of a compact Kähler manifold M onto a smooth compact complex curve C. Consider a flat vector bundle L on C of complex rank 1 with unitary structure group and let $E = f^*L$ be the pullback of this bundle to M. Let $\omega \in \Omega^1(E)$ be a holomorphic 1-form with values in E (by the Hodge decomposition ω is d-closed).

Lemma 3.1 *Assume that $\omega|_{V_z} = 0$ for the fibre $V_z := f^{-1}(z)$ over a regular value $z \in C$. Then $\omega|_V = 0$ for any fibre V of f.*

Remark 3.2 If V is not smooth the lemma asserts that ω equals 0 on the smooth part of V.

Proof. Denote by S the set of non-regular values of f. Then $S \subset C$ consists of a finite number of points. First consider fibres of f over regular values. By Sard's theorem, $f : M \setminus f^{-1}(S) \longrightarrow C \setminus S$ is a fibre bundle with connected fibres. According to our assumption there is a fibre V_z of this bundle such that $\omega|_{V_z} = 0$. Then $\omega|_V = 0$ for any fibre V of $f|_{M \setminus f^{-1}(S)}$. In fact, for any fibre V there is an open neighbourhood O_V of V diffeomorphic to $\mathbb{R}^2 \times V$ such that $E|_{O_V}$ is the trivial flat vector bundle (here we have used the fact that $E|_V$ is trivial because it is pullback of a bundle defined over a point). Any d-closed holomorphic 1-form defined on O_V that vanishes on V is d-exact, and so it vanishes on each fibre contained in O_V. Starting with a tubular neighbourhood O_{V_z} and taking into account that $\omega|_{O_{V_z}}$ can be considered as a d-closed holomorphic 1-form (because $E|_{O_{V_z}}$ is trivial) we obtain that $\omega|_V = 0$ for any fibre $V \subset O_{V_z}$. Then the required statement follows by induction if we cover $M \setminus f^{-1}(S)$ by open tubular neighbourhoods of fibres of f and use the fact that $C \setminus S$ is connected.

Consider now fibres over the singular part S. Let $O_x \subset C$ be the neighbourhood of a point $x \in S$ such that $O_x \setminus S = O_x \setminus \{x\}$ is biholomorphic to $\mathbb{D} \setminus \{0\}$. In particular, $\pi_1(O_x \setminus S) = \mathbb{Z}$. Without loss of generality we may assume that $V = f^{-1}(x)$ is a deformation retract of an open neighbourhood U_x of V and $W_x := f^{-1}(O_x) \subset U_x$ (such U_x exists, e.g., by the triangulation theorem of the pair (M, V), see [L, Th.2]). Since $E|_V$ is trivial, the bundle $E|_{U_x}$ is also trivial. Therefore we can regard ω as a d-closed holomorphic 1-form on U_x. Moreover, we already have proved that $\omega|_F = 0$ for any fibre $F \subset W_x \setminus f^{-1}(S)$. So ω equals 0 on each fibre of the fibration $f : W_x \setminus f^{-1}(S) \longrightarrow O_x \setminus S$. This implies that there is a d-closed holomorphic 1-form ω_1 on $O_x \setminus S$ such that $\omega = f^*(\omega_1)$. Assume now that $\omega|_V \neq 0$. Then integration of $\omega|_{W_x}$ along paths determines a non-trivial homomorphism $h_{\omega,x} : \pi_1(W_x) \longrightarrow \mathbb{C}$ whose image is isomorphic to $\mathbb{Z}$. In fact, since embedding $W_x \setminus f^{-1}(S) \hookrightarrow W_x$ induces a surjective homomorphism of fundamental groups, we can integrate ω by paths contained in $W_x \setminus f^{-1}(S)$ obtaining the same image in $\mathbb{C}$. Further, for any path $\gamma \subset W_x \setminus f^{-1}(S)$ we have $\int_\gamma \omega = \int_{\gamma_1} \omega_1$, where γ_1 is a path in $O_x \setminus S$ representing the element $f_*(\gamma) \in \pi_1(O_x \setminus S)$. But $\pi_1(O_x \setminus S) = \mathbb{Z}$ and therefore $h_{\omega,x}(\pi_1(W_x)) \cong \mathbb{Z}$. Recall that any path in W_x is homotopically equivalent inside U_x to a path contained in V and so $\omega|_V$ determines a homomorphism h of $\Gamma := (\pi_1(V)/D\pi_1(V))/_{torsion}$ into $\mathbb{C}$ with image isomorphic to $\mathbb{Z}$. Without loss of generality we may assume that V is smooth (for otherwise, we apply the arguments below to each irreducible component of a desingularization of V). Integrating holomorphic 1-forms on V along paths we embed Γ into some $\mathbb{C}^k$ as a lattice of rank $2k$. Then there is a linear holomorphic functional f_ω on $\mathbb{C}^k$ such that $h = f_\omega|_\Gamma$. Since $rk(h(\Gamma)) = 1$, f_ω equals 0 on a subgroup $H \subset \Gamma$ isomorphic to $\mathbb{Z}^{2k-1}$. In particular, $f_\omega = 0$ on the vector space $R = span(H)$ of real dimension $2k-1$. But $Ker(f_\omega)$ is a complex vector space. So $f_\omega = 0$ on $\mathbb{C}^k$ which implies that $\omega|_V = 0$. This contradicts our assumption and proves the required statement for fibres over the points of S.

The lemma is proved. $\square$

3.2. Let $f : M \longrightarrow C$ be a surjective holomorphic map with connected fibres of a compact Kähler manifold M onto a smooth compact complex curve C. Then the induced homomorphism $f_* : \pi_1(M) \longrightarrow \pi_1(C)$ is surjective. Denote by $T_2 \subset T_2(\mathbb{C})$ the Lie group of matrices of the form

$$\begin{pmatrix} a & b \\ 0 & 1 \end{pmatrix} \qquad (a \in \mathbb{C}^*,\ b \in \mathbb{C}).$$

For a homomorphism $\rho \in Hom(\pi_1(M), T_2)$ we let $\rho_a \in Hom(\pi_1(M), \mathbb{C}^*)$ denote the upper diagonal character of ρ.

Proposition 3.3 *Let $\rho \in Hom(\pi_1(M), T_2)$ be such that ρ_a is a non-unitary character. Assume that $\rho_a|_{Ker(f_*)}$ is trivial. Then $\rho|_{Ker(f_*)}$ is also trivial.*

Proof. Let $p_t : M_t \longrightarrow M$ be the Galois covering with transformation group $Tor(\pi_1(M)/D\pi_1(M))$. Then $\rho|_{\pi_1(M_t)}$ determines a C^∞-trivial complex rank-2 flat vector bundle on M_t, because $\rho_a|_{\pi_1(M_t)} = \exp(\rho'|_{\pi_1(M_t)})$ for some $\rho' \in Hom(\pi_1(M), \mathbb{C})$. In particular, $\rho|_{\pi_1(M_t)}$ can be defined by a flat connection

$$\Omega = \begin{pmatrix} \omega & \eta \\ 0 & 0 \end{pmatrix}; \qquad d\Omega - \Omega \wedge \Omega = 0$$

on $M_t \times \mathbb{C}^2$ where ω is a d-harmonic 1-form on M_t lifted from M. Moreover, since $\rho_a|_{Ker(f_*)}$ is trivial, ω is the pullback by $f \circ p_t$ of a d-harmonic 1-form defined on C. Let $\omega = \omega_1 + \omega_2$ be the type decomposition of ω into the sum of holomorphic and antiholomorphic 1-forms lifted from C. Denote by E_ρ the rank-1 flat vector bundle on M_t with unitary structure group constructed by the flat connection $\omega_2 - \overline{\omega_2}$ and by $E_0 = M_t \times \mathbb{C}$ the trivial flat vector bundle. Observe that E_ρ is pullback of a flat bundle on C. In particular, $E_\rho|_V$ is trivial for any fibre V of $f \circ p_t$. According to the results of Sections 2.1 and 2.2 the equivalence class of $\rho|_{\pi_1(M_t)}$ is determined by a d-harmonic 1-form θ with values in $End(E_\rho \oplus E_0)$ satisfying $\theta \wedge \theta$ represents 0 in $H^2(M_t, End(E_\rho \oplus E_0))$. More precisely,

$$\theta = \begin{pmatrix} \omega_1 & \eta' \\ 0 & 0 \end{pmatrix}$$

where η' is a d-harmonic 1-form with values in E_ρ satisfying $\omega_1 \wedge \eta'$ represents 0 in $H^2(M_t, E_\rho)$. Let $M_t \xrightarrow{g_1} M_1 \xrightarrow{g_2} C$ be the Stein factorization of $f \circ p_t$. Here g_1 is a morphism with connected fibres onto a smooth curve M_1 and g_2 is a finite morphism. Then for any fibre $V \hookrightarrow M_t$ of g_1 we have that $p_t|_V : V \longrightarrow p_t(V)$ is a regular covering of the fibre $p_t(V)$ of f with a finite abelian transformation group.

Lemma 3.4 *$\eta'|_V = 0$ for fibres over regular values of g_1.*

Based on this statement we, first, finish the proof of the proposition and then will prove the lemma. From the lemma and Lemma 3.1 it follows that $\eta'|_W = 0$ for any fibre $i : W \hookrightarrow M_t$ of g_1. Thus $\rho|_{\pi_1(M_t)}$ is trivial on $i_*(\pi_1(W)) \subset \pi_1(M_t)$. But $\pi_1(W)$ is a subgroup of a finite index in $\pi_1(p_t(W))$ and the image of $\rho|_{j_*(\pi_1(p_t(W)))}$ consists of unipotent matrices by the assumption of the proposition. (Here $j : p_t(W) \hookrightarrow M$ is embedding.) Therefore $\rho|_{j_*(\pi_1(p_t(W)))}$ is trivial for any W. Denote by E the flat vector bundle on M associated to ρ. Then we have proved that $E|_W$ is trivial for any fibre W of f. Further, let $(U_i)_{i \in I}$ be an open covering of C such that $W_i := f^{-1}(U_i)$ is an open neighbourhood of a fibre $V_i = f^{-1}(x_i)$, $x_i \in U_i$, and W_i is deformable onto V_i. Since $E|_{V_i}$ is trivial, $E|_{W_i}$ is also trivial. In particular, E is defined by a locally constant cocycle $\{c_{ij}\}$ defined on the covering $(W_i)_{i \in I}$. But then $\{c_{ij}\}$ is the pullback of a cocycle defined on $(U_i)_{i \in I}$ because the fibres of f are connected. This cocycle determines a bundle E' on C such that $f^*E' = E$. Then ρ is the pullback of a homomorphism $\rho' \in Hom(\pi_1(C), T_2)$ constructed by E'.

This completes the proof of the proposition modulo Lemma 3.4.

Proof of Lemma 3.4. Observe that $\omega_1 \neq 0$ in the above definition of θ because ρ_a is non-unitary. We also regard $\eta' \neq 0$, for otherwise, the image of ρ consists of diagonal matrices and the required statement is trivial. Let $i : V \hookrightarrow M_t$ be a fibre over a regular value of g_1. For any $\lambda \in \mathbb{C}^*$ consider the form

$$\theta_\lambda = \begin{pmatrix} \lambda\omega_1 & \eta' \\ 0 & 0 \end{pmatrix}.$$

Clearly, $\theta_\lambda \wedge \theta_\lambda$ represents 0 in $H^2(M_t, End(E_\rho \oplus E_0))$ and thus, according to the results of Section 2.2, it determines a representation $\rho_\lambda \in Hom(\pi_1(M_t), T_2)$ with the upper diagonal character $\rho_{\lambda a}$ defined by the flat connection $\lambda\omega_1 + \omega_2$. In particular, the family $\{\rho_{\lambda a}\}$ contains infinitely many different characters. Assume that $\rho|_{i_*(\pi_1(V))}$ is not trivial. Then it can be determined by the restriction $\theta|_{i(V)}$. But according to our assumption $\omega_1|_{i(V)} = 0$ and $E_\rho|_{i(V)}$ is a trivial flat vector bundle. Thus $\rho|_{i_*(\pi_1(V))}$ is defined by $\eta'|_{i(V)} \in H^1(i(V), \mathbb{C})$. Let $\psi \in Hom(i(V), \mathbb{C})$ be a homomorphism obtained by integration of $\eta'|_{i(V)}$ along paths generating $\pi_1(i(V))$. Then

$$\rho|_{i_*(\pi_1(V))} = \begin{pmatrix} 1 & \psi \\ 0 & 1 \end{pmatrix}.$$

We also obtain that $\rho_\lambda|_{i_*(\pi_1(V))} = \rho|_{i_*(\pi_1(V))}$ and so it is not trivial for any $\lambda \in \mathbb{C}^*$. We will prove now that the family $\{\rho_{\lambda a}\}$ contains finitely many different characters. This contradicts our assumption and shows that $\rho|_{i_*(\pi_1(V))}$ is trivial and so $\eta'|_V = 0$.

Let $H := i_*(\pi_1(V))$ be a normal subgroup of $\pi_1(M_t)$ (the normality follows from the exact homotopy sequence for the bundle defined over regular values of g_1). Denote by H_{ab} the quotient H/DH. Then H_{ab} is an abelian group of a finite rank. Moreover, H_{ab} is a normal subgroup of $K := \pi_1(M_t)/DH$ and the group $K_1 := K/H_{ab}$ is finitely generated. Note that ρ_λ induces a homomorphism $\widehat{\rho}_\lambda : K \longrightarrow T_2$ for any $\lambda \in \mathbb{C}^*$ with the same image as for ρ_λ because the image of $\rho_\lambda|_H$ is abelian. In particular, $\widehat{\rho}_\lambda(H_{ab})$ is a normal subgroup of $\widehat{\rho}(K)$. Since by our assumption $\widehat{\rho}_\lambda(H_{ab})$ is a non-trivial subgroup of unipotent matrices, from the identity

$$\begin{pmatrix} c & b \\ 0 & 1 \end{pmatrix} \cdot \begin{pmatrix} 1 & v \\ 0 & 1 \end{pmatrix} \cdot \begin{pmatrix} c & b \\ 0 & 1 \end{pmatrix}^{-1} = \begin{pmatrix} 1 & c \cdot v \\ 0 & 1 \end{pmatrix} \qquad (c \in \mathbb{C}^*,\ b,\ v \in \mathbb{C})$$

it follows that the action of $\widehat{\rho}_\lambda(K)$ on $\widehat{\rho}_\lambda(H_{ab})$ by conjugation is defined by multiplication of non-diagonal elements of $\widehat{\rho}_\lambda(H_{ab})$ by elements of $\rho_{\lambda a}(\pi_1(M_t))$. Note also that $Tor(H_{ab})$ belongs to $Ker(\widehat{\rho}_\lambda)$ for any λ. Thus if $H' \cong \mathbb{Z}^s$ is a maximal free abelian subgroup of H_{ab} then $\rho_{\lambda a}(\pi_1(M_t))$ consists of eigen values of matrices from $SL_s(\mathbb{Z})$ obtained by the natural action (by conjugation) of K_1 on H'. Since K_1 is finitely generated, the number of different $\rho_{\lambda a}$ is finite which is false.

This contradiction completes the proof of the lemma. □

3.3. Let $f : M \longrightarrow C$ be a holomorphic map with connected fibres of a compact Kähler manifold M onto a smooth compact complex curve C of genus $g \geq 1$. Then $\pi_1(M)$ is defined by the exact sequence

$$\{e\} \longrightarrow Ker(f_*) \longrightarrow \pi_1(M) \longrightarrow \pi_1(C) \longrightarrow \{e\}\ .$$

Moreover, $Ker(f_*)$ is a finitely generated group. Let $G \subset Ker(f_*)$ be a normal subgroup of $\pi_1(M)$. Then the quotient group $R := \pi_1(M)/G$ is defined by the sequence

$$\{e\} \longrightarrow Ker(f_*)/G \longrightarrow R \longrightarrow \pi_1(C) \longrightarrow \{e\} .$$

Assume that

(1) $H := Ker(f_*)/G$ is a free abelian group of rank $k \geq 1$;
(2) the natural action of $D\pi_1(C)$ on H is trivial.

From (2) it follows that the action s of $\pi_1(C)$ on H determines an action of $\mathbb{Z}^{2g} \cong \pi_1(C)/D\pi_1(C)$ on H. Identifying H with $\mathbb{Z}^k$ we can think of H as a subgroup (lattice) of $Z^k \otimes \mathbb{C} = \mathbb{C}^k$. Then s determines a representation $s' : \pi_1(C) \longrightarrow GL_k(\mathbb{C})$ such that $Ker(s') \subset D\pi_1(C)$ and $s'(g)|_H = s(g)$ for any $g \in \pi_1(C)$. Since s' descends to a representation $\mathbb{Z}^{2g} \longrightarrow GL_k(\mathbb{C})$, it admits a decomposition $s' = \oplus_{j=1}^m s_j$ where s_j is equivalent to a nilpotent representation $\pi_1(C) \longrightarrow T_{k_i}(\mathbb{C})$ with a diagonal character ρ_j. Here $\sum_{j=1}^m k_j = k$.

Proposition 3.5 *All characters ρ_j are torsion.*

Proof. By definition, R is an extension of $\pi_1(C)$ by H. It is well known, see, e.g. [G, Ch.I, Sec.6], that the class of extensions equivalent to R is uniquely defined by an element $c \in H^2(\pi_1(C), H)$ where the cohomology is defined by the action s of $\pi_1(C)$ on H. Let $f \in Z^2(\pi_1(C), H)$ be a cocycle determining c. Then one can define a representative of the equivalence class of extensions as the direct product $H \times \pi_1(C)$ with multiplication

$$(h_1, g_1) \cdot (h_2, g_2) = (h_1 + s(g_1)(h_2) + f(g_1, g_2), g_1 \cdot g_2);$$

$$h_1, h_2 \in H,\ g_1, g_2 \in \pi_1(C) .$$

The natural embedding $H \hookrightarrow \mathbb{Z}^k \otimes \mathbb{C}(= \mathbb{C}^k)$ determines an embedding i of R into the group R' defined as $\mathbb{C}^k \times \pi_1(C)$ with multiplication

$$(v_1, g_1) \cdot (v_2, g_2) = (v_1 + s'(g_1)(v_2) + f(g_1, g_2), g_1 \cdot g_2);$$

$$f \in Z^2(\pi_1(C), H),\ v_1, v_2 \in \mathbb{C}^k,\ g_1, g_2 \in \pi_1(C) .$$

Here we regard f as an element of $Z^2(\pi_1(C), \mathbb{C}^k)$ defined by the action s'. From the decomposition $s' = \oplus_{j=1}^m s_j$ it follows that there is an invariant $\pi_1(C)$-submodule $V_j \subset \mathbb{C}^k$ of $dim_{\mathbb{C}} V_j = k - 1$ such that $W_j = \mathbb{C}^k/V_j$ is a one-dimensional $\pi_1(C)$-module and the action of $\pi_1(C)$ on W_j is defined as multiplication by the character ρ_j. Then, by definition, V_j is a normal subgroup of R' and the the quotient group $R_j = R'/V_j$ is defined by the sequence

$$\{e\} \longrightarrow \mathbb{C} \longrightarrow R_j \longrightarrow \pi_1(C) \longrightarrow \{e\} .$$

Here the action of $\pi_1(C)$ on $\mathbb{C}$ is multiplication by the character ρ_j. Further, the equivalence class of extensions isomorphic to R_j is defined by an element $c_j \in H^2(\pi_1(C), \mathbb{C})$ (the cohomology is defined by the above action of $\pi_1(\mathbb{C})$ on $\mathbb{C}$). We will assume that the character ρ_j is non-trivial (for otherwise, ρ_j is clearly torsion). Let us denote by t_j the composite homomorphism $\pi_1(M) \longrightarrow R \xrightarrow{i} R' \longrightarrow R_j$.

Let E_{ρ_j} be a complex rank-1 flat vector bundle on C constructed by $\rho_j \in Hom(\pi_1(C), \mathbb{C}^*)$. Since C is a $K(\pi_1(C), 1)$-space, there is a natural isomorphism of the above group $H^2(\pi_1(C), \mathbb{C})$ and the Čech cohomology group $H^2(C, \mathbf{E}_{\rho_j})$ of the sheaf of locally constant sections of E_{ρ_j}, see e.g. [M, Ch.1, Complement to Sec.2]. But each flat vector bundle on C is C^∞-trivial and each homomorphism from $Hom(\pi_1(C), \mathbb{C}^*)$ can be continuously deformed inside of $Hom(\pi_1(C), \mathbb{C}^*)$ to the trivial homomorphism. Therefore by the index theorem

$$dim_{\mathbb{C}} H^0(C, \mathbf{E}_{\rho_\mathbf{j}}) - dim_{\mathbb{C}} H^1(C, \mathbf{E}_{\rho_\mathbf{j}}) + dim_{\mathbb{C}} H^2(C, \mathbf{E}_{\rho_\mathbf{j}}) = \chi(\pi_1(C)) = 2 - 2g \ .$$

Note that $H^0(C, \mathbf{E}_{\rho_\mathbf{j}}) = 0$ because ρ_j is non-trivial. Furthermore, $H^1(C, \mathbf{E}_{\rho_\mathbf{j}})$ is in a one-to-one correspondence with the set of non-equivalent representations $\rho : \pi_1(C) \longrightarrow T_2 \subset T_2(\mathbb{C})$ with the upper diagonal character ρ_j. Using the identity $\prod_{i=1}^{g} [e_i, e_{g+i}] = e$ for generators $e_1, .., e_{2g} \in \pi_1(C)$ we easily obtain that $dim_{\mathbb{C}} H^1(C, \mathbf{E}_{\rho_\mathbf{j}}) = 2g - 2$. Thus we have $H^2(C, \mathbf{E}_{\rho_\mathbf{j}}) = 0$. This shows that $H^2(\pi_1(C), \mathbb{C}) = 0$ and R_j is isomorphic to the semidirect product of $\mathbb{C}$ and $\pi_1(C)$, i.e., $R_j = \mathbb{C} \times \pi_1(C)$ with multiplication

$$(v_1, g_1) \cdot (v_2, g_2) = (v_1 + \rho_j(g_1) \cdot v_2, g_1 \cdot g_2), \quad v_1, v_2 \in \mathbb{C},\ g_1, g_2 \in \pi_1(C) \ .$$

Let us determine a map ϕ_j of R_j to T_2 by the formula

$$\phi_j(v, g) = \begin{pmatrix} \rho_j(g) & v \\ 0 & 1 \end{pmatrix}$$

Obviously, ϕ_j is a correctly defined homomorphism with upper diagonal character ρ_j. Hence $\phi_j \circ t_j : \pi_1(M) \longrightarrow T_2(\mathbb{C})$ is a homomorphism which is non-trivial on $Ker(f_*)$ by its definition. Now Proposition 3.3 implies that ρ_j is a unitary character. Therefore we have proved that all characters of the action of $\pi_1(C)$ on H are unitary. This means that each element of the above action is defined by a matrix from $SL_k(\mathbb{Z})$ with unitary eigen values. Applying now the theorem on the units of a ring of integers of an algebraic field (see, e.g. [BS, p.105,Th.2]) we obtain that each ρ_j is a torsion character. □

Using Proposition 3.5 we prove the following statement.

Let $f : M \longrightarrow C$ be a holomorphic map with connected fibres of a compact Kähler manifold M onto a smooth compact complex curve C of genus $g \geq 1$.

Proposition 3.6 *Assume that $\rho \in Hom(\pi_1(M), T_2)$ and that the upper diagonal character ρ_a is pullback of a character from $Hom(\pi_1(C), \mathbb{C}^*)$. Assume also that $\rho|_{Ker(f_*)}$ is non-trivial. Then ρ_a is torsion. In addition, the number of ρ_a satisfying these assumptions is finite.*

Proof. First, from the assumption of the proposition it follows that ρ maps $Ker(f_*)$ to the abelian group $N := DT_2$ of unipotent matrices in $T_2(\mathbb{C})$. Let G be the intersection of kernels of all homomorphisms ρ satisfying the conditions of the proposition.

Then G is a normal subgroup of $\pi_1(M)$ and $D(Ker(f_*))$ is contained in G. Set $G_1 := Ker(f_*) \cap G$ and let $R := \pi_1(M)/G_1$, $H := Ker(f_*)/G_1$. Then R is defined by the exact sequence

$$\{e\} \longrightarrow H \longrightarrow R \longrightarrow \pi_1(C) \longrightarrow \{e\}\ .$$

Observe that by the definition of G_1 and conditions of the proposition, H is a non-trivial finitely generated free abelian group. Let us check that $D\pi_1(C)$ acts trivially on H. In fact, let $a \in D\pi_1(M)$, $b \in Ker(f_*)$. Then $\rho([a,b]) = [\rho(a), \rho(b)] \subset DN = \{e\}$ for any homomorphism ρ satisfying conditions of the proposition. Thus $[a,b] \in G$. Moreover, $[a,b] \in Ker(f_*)$ because this is a normal subgroup. Thus $[a,b] \in G_1$ and so its image in R is trivial. This implies the required statement. Further, according to Proposition 3.5 all characters of the action s of $\pi_1(C)$ on H are torsion. Consider now a homomorphism ρ satisfying assumptions of the proposition. Then clearly ρ determines a homomorphism $\rho' : R \longrightarrow T_2$ with the same image. Since according to our assumption $\rho|_{Ker(f_*)}$ is non-trivial, $\rho'|_H$ is also non-trivial. Thus $\rho'(H)$ is a non-trivial normal subgroup of $\rho'(R)$. But $\rho'(H)$ is a subgroup of unipotent matrices. Hence, the action of $\rho'(R)$ on $\rho'(H)$ by conjugation is defined by multiplication of the non-diagonal elements of $\rho'(R)$ by elements of $\rho_a(\pi_1(M))$. So $\rho_a(\pi_1(M))$ consists of eigen values of invertible integer matrices obtained by the natural action s of $\pi_1(C)$ on H. This implies that ρ_a is torsion because as we have proved, all characters of s are torsion. This argument also shows that the number of ρ_a satisfying conditions of the proposition is finite. $\square$

3.4. Proof of Theorem 1.9. Let V_ρ be a flat vector bundle on M associated to $\rho \in Hom(\pi_1(M), T_n(\mathbb{C}))$ satisfying the assumptions of the theorem. Since the homomorphism $d(\rho) \in Hom(\pi_1(M), D_n(\mathbb{C}))$ is the pullback of a homomorphism from $Hom(\pi_1(C), D_n(\mathbb{C}))$, V_ρ is C^∞-trivial. Let V_i be a flat vector bundle associated to the homomorphism ρ''_{iu}. Then $V_\xi = \oplus_{i=1}^n V_i$ is a flat vector bundle on C associated to $\xi := diag[\rho''_{1u}, ..., \rho''_{nu}] \in Hom(\pi_1(C), D_n^u)$. According to the results of Sections 2.1 and 2.2, the equivalence class of ρ is uniquely defined by an upper triangular $End(f^*V_\xi)$-valued d-harmonic 1-form α with a nilpotent (0,1)-component such that $\alpha \wedge \alpha$ represents 0 in $H^2(M, End(f^*V_\xi))$. Moreover, according to (1) $diag(\alpha_1)$ of the (1,0)-component α_1 of α is the pullback of a harmonic 1-form on C. Further, $\alpha = \sum_{i,j=1}^n \alpha_{ij}$ where α_{ij} is a d- harmonic 1-form with values in $f^*(V_i^* \otimes V_j)$. Any pair $(f^*(V_i^* \otimes V_j), \alpha_{ij})$ determines a homomorphism $\rho_{ij} \in Hom(\pi_1(M), T_2)$ with upper diagonal character $f^*(\rho''_{iu} \otimes (\rho''_{ju})^{-1})$. Since by assumption (2) the latter character is not torsion, Proposition 3.5 implies that ρ_{ij} is the pullback of a representation $\pi_1(C) \longrightarrow T_2$. In particular, there is a d- harmonic 1-form β_{ij} with values in $V_i^* \otimes V_j$ such that $f^*(\beta_{ij}) = \alpha_{ij}$. Thus α is the pullback of an upper triangular $End(V_\xi)$-valued harmonic 1-form $\beta = \sum_{i,j=1}^n \beta_{ij}$ with a nilpotent (0,1)-component. Now Example 2.1 1 says that the pair (V_ξ, β) determines a representation $\widetilde{\rho} \in Hom(\pi_1(C), T_n(\mathbb{C}))$. Finally, our construction and results of Section 2.2 give $f^*(\widetilde{\rho}) = \rho$. $\square$

3.5. Proof of Theorem 1.11. Let M be a compact Kähler manifold with fundamental group F. According to the assumption of the theorem, F admits a surjective homomorphism p onto a fundamental group G of a compact Riemann surface of genus $g \geq 2$. (Here p is the composition $F \xrightarrow{j} G_1 \longrightarrow G_1/\mathbb{Z}^k = G$.) Then by the Siu-Beauville theorem (see, e.g. [ABCKT, Th.2.11]) there are a holomorphic map $f : M \longrightarrow C$ with connected fibres onto a compact complex curve C of genus $g' \geq g$ and a homomorphism $h : \pi_1(C) \longrightarrow G$ such that $p = h \circ f_*$. Let $E := f_*(j^{-1}(\mathbb{Z}^k)) \subset \pi_1(C)$ be a normal subgroup. Assume first that $\pi_1(C)/E$ is finite. Then we have $h(E)$ is a normal subgroup of a finite index in G. But $h(E) = p(j^{-1}(\mathbb{Z}^k)) = \{e\}$ which is wrong. Thus $\pi_1(C)/E$ is infinite and so E is a free group. According to the assumption of the theorem the number of generators r of E is finite. Let $S \longrightarrow C$ be a regular covering corresponding to $\pi_1(C)/E$. If $r \geq 2$ then the group $Iso(S)$ of isometries of S is finite and since $\pi_1(C)/E$ is infinite we have $r \leq 1$. If $r = 1$, any discrete subgroup of $Iso(S)$ is virtually cyclic and in particular does not act cocompactly on S. Thus $r = 0$ which means that $j^{-1}(\mathbb{Z}^k) \subset Ker(f_*)$, $\pi_1(C) \cong G$ and h is the identity homomorphism. In particular, $K \subset Ker(f_*)$ satisfies conditions (1), (2) of Proposition 3.5. This implies that all characters of the action of G on $\mathbb{Z}^k$ are torsion. □

4. Proofs of Theorems 1.1, 1.4 and 1.6.

Sections 4.1 and 4.2 below contain the proof of Theorem 1.1.

4.1. First we prove a two-dimensional version of the required result.

Let $T_2^u \subset T_2(\subset T_2(\mathbb{C}))$ be the subgroup of matrices with unitary upper diagonal elements. As before, $\rho_a \in Hom(\pi_1(M), U_1)$ denotes the upper diagonal character of $\rho \in Hom(\pi_1(M), T_2^u)$. Recall that X_a stands for the normalization of the image of the Albanese map $\alpha_X : X \longrightarrow Alb(X)$ and $\alpha_X^n : X \longrightarrow X_a$ denotes a holomorphic map which covers α_X.

Lemma 4.1 *There is a finite Galois covering $r : M_H \longrightarrow M$ with an abelian Galois group $H := H(M)$ such that for any $\rho \in Hom(\pi_1(M), T_2^u)$ there is $\widetilde{\rho} \in Hom(\pi_1(M_{Ha}), T_2^u)$ so that $\rho \circ r_* = \widetilde{\rho} \circ \alpha^n_{M_H *}$.*

Let us introduce some definitions. Given a character $\xi \in Hom(\pi_1(M), \mathbb{C}^*)$, let $\mathbb{C}_\xi$ denote the associated $\pi_1(M)$-module. We define $\Sigma^1(\pi_1(M))$ to be the set of characters $\xi \in Hom(\pi_1(M), \mathbb{C}^*)$ such that $H^1(\pi_1(M), \mathbb{C}_\xi)$ is nonzero.

Theorem ([BSC]). *There is a finite number of surjective holomorphic maps with connected fibres onto smooth compact complex curves $f_i : M \longrightarrow C_i$ and torsion characters $\rho_i, \rho'_j \in Hom(\pi_1(M), \mathbb{C}^*)$ such that*

$$\Sigma^1(\pi_1(M)) = \bigcup_i \rho_i f_i^*(Hom(\pi_1(C_i), \mathbb{C}^*)) \cup \bigcup_j \{\rho'_j\} \ .$$

Remark 4.2 The structure of the positive dimensional components of $\Sigma^1(\pi_1(M))$ was discovered by Beauville [Be]. In [S1] Simpson proved that the set of zero dimensional components of $\Sigma^1(\pi_1(M))$ consists of torsion characters provided M is a smooth projective manifold. Finally, Campana [C1] established this fact for an arbitrary compact Kähler manifold by reducing the question to Simpson's theorem (the projective case).

Proof of Lemma 4.1. We represent $Hom(\pi_1(M), T_2^u)$ as $R_1 \cup R_2 \cup R_3 \cup R_4$ with R_i to be defined below and prove the lemma for any component of the decomposition.

It is well-known that the equivalence class of $\rho \in Hom(\pi_1(M), T_2^u)$ is defined by an element of $H^1(\pi_1(M), \mathbb{C}_{\rho_a})$. Let R_1 consist of homomorphisms ρ such that $H^1(\pi_1(M), \mathbb{C}_{\rho_a}) = 0$. Then ρ is equivalent to a representation in the group of diagonal 2×2 matrices. Thus if $p_t : M_t \longrightarrow M$ is the Galois covering with the Galois group $Tor(\pi_1(M)/D\pi_1(M))$ then obviously $\rho|_{\pi_1(M_t)}$ is the pullback by α_{M_t} of a homomorphism from $Hom(\pi_1(Alb(M_t)), T_2^u)$.

Assume now that $H^1(\pi_1(M), \mathbb{C}_{\rho_a}) \neq 0$, or equivalently, $\rho_a \in \Sigma^1(\pi_1(M))$. Let K be intersection of the kernels of all characters ρ_i and ρ'_j from the BSC theorem. Then K is a normal subgroup of $\pi_1(M)$ and the quotient group $G := \pi_1(M)/K$ is finite abelian. Let $p : M_G \longrightarrow M$ be the Galois covering with the Galois group G. We set $\rho'_a := \rho_a|_{\pi_1(M_G)}$. Then ρ'_a is the upper diagonal character of $\rho' := \rho|_{\pi_1(M_G)}$. Further, according to the BSC theorem, there are two possibilities:
(1) ρ'_a is trivial;
(2) there are a holomorphic surjective map f with connected fibres onto a smooth curve C and $\rho_1 \in Hom(\pi_1(C), \mathbb{C}^*)$ such that $\rho'_a = \rho_1 \circ (f \circ p)_*$.
We will denote by R_2 the set of $\rho \in Hom(\pi_1(M), T_2^u)$ satisfying (1). Then the image of $\rho|_{\pi_1(M_G)}$ for $\rho \in R_2$ is abelian and consists of unipotent matrices. Therefore $\rho|_{\pi_1(M_G)}$ is the pullback by α_{M_G} of a homomorphism from $Hom(\pi_1(\alpha_{M_G}(M_G)), T_2^u)$.

Consider now case (2). Let $M_G \xrightarrow{g_1} C_1 \xrightarrow{g_2} C$ be the Stein factorization of $f \circ p$. Here g_1 is a morphism with connected fibres onto a smooth curve C_1 and g_2 is a finite morphism. Then ρ'_a is the pullback by g_1 of $g_2^*(\rho_1) \in Hom(\pi_1(C_1), \mathbb{C}^*)$. Denote by $R_3(f)$ the set of $\rho \in Hom(\pi_1(M), T_2^u)$ for which $\rho'|_{Ker(g_{1*})}$ is non-trivial and let $R_3 = \cup_i R_3(f_i)$ where the union is taken over all maps f_i from the BSC theorem. According to Proposition 3.6, ρ'_a for $\rho \in R_3(f)$ is torsion and the number of such characters is finite. This implies that ρ_a is also a torsion character. Moreover, if $n(\rho)$ is the number of distinct ρ_a for $\rho \in R_3(f)$ then $\sum_i \sum_{\rho \in R_3(f_i)} n(\rho) < \infty$.

Finally, denote by $R_4(f)$ the set of homomorphisms ρ for which $\rho'|_{Ker(g_{1*})}$ is trivial and let $R_4 = \cup_i R_4(f_i)$. Then for any $\rho \in R_4(f)$ there is $\xi \in Hom(\pi_1(C_1), T_2^u)$ such that $\rho' = \xi \circ g_{1*}$.

Let F be the intersection of the kernels of all characters ρ_a for all $\rho \in R_3$ and $F_1 := F \cap K \cap \pi_1(M_t)$. Clearly F_1 is a normal subgroup of $\pi_1(M)$ and $H := \pi_1(M)/F_1$ is finite abelian. Let $r : M_H \longrightarrow M$ be the Galois covering with Galois group H. Then there is a covering map $r_1 : M_H \longrightarrow M_G$ such that $r = p \circ r_1$. Now for $\rho \in R_3(f) \cup R_4(f)$ let $g_1 : M_G \longrightarrow C_1$ be as above and $M_H \xrightarrow{h_1} C_2 \xrightarrow{h_2} C_1$ be the Stein factorization of $g_1 \circ r_1$. Here h_1 is a morphism with connected fibres onto a smooth curve C_2 and h_2 is a finite morphism. Then $\rho''_a := \rho_a|_{\pi_1(M_H)}$ is the pullback by h_1 of $(g_2 \circ h_2)^*(\rho_1) \in Hom(\pi_1(C_2), \mathbb{C}^*)$ and ρ''_a is the upper diagonal character of $\rho'' := \rho|_{\pi_1(M_H)}$. In particular, if $\rho \in R_4(f)$ then ρ'' is pullback by h_1 of $\xi \circ h_{2*} \in Hom(\pi_1(C_2), T_2^u)$. Since C_2 is a smooth curve, the Albanese map $\alpha_{C_2} : C_2 \longrightarrow Alb(C_2)$ is an embedding. Below we identify C_2 with $\alpha_{C_2}(C_2)$. Then there is a holomorphic surjective map $a_{C_2} : Alb(M_H) \longrightarrow Alb(C_2)$ such that $h_1 = a_{C_2} \circ \alpha_{M_H}$. Thus if $n : M_{Ha} \longrightarrow \alpha_{M_H}(M_H)$ is normalization, then ρ'' is the pullback by $\alpha^n_{M_H}$ of $\xi \circ (h_2 \circ a_{C_2} \circ n)_* \in Hom(\pi_1(M_{Ha}), T_2^u)$.

Assume now that $\rho \in R_3(f)$. Then ρ''_a is trivial and the image of ρ'' consists of unipotent matrices (and so is abelian). This implies that ρ'' is pullback of a homomorphism from M_{Ha} because any such ρ'' is defined by integration of a harmonic 1-form on M_H.

So we have considered all possible implications for $\rho \in Hom(\pi_1(M), T_2^u)$ and have proved Lemma 4.1 with the above defined H. $\square$

Remark 4.3 1. Let $\widetilde{\rho} \in Hom(\pi_1(M_{Ha}), T_2^u)$ be a homomorphism from Lemma 4.1, that is $\rho \circ r_* = \widetilde{\rho} \circ \alpha^n_{M_{H*}}$, and $\widetilde{\rho}_a$ be the upper diagonal character of $\widetilde{\rho}$. Then the arguments of the lemma show that $\widetilde{\rho}_a = \exp(\eta)$ for some $\eta \in Hom(\pi_1(M_{Ha}), \mathbb{C})$. In particular, $\widetilde{\rho}_a$ is uniquely defined by ρ_a. Indeed, by definition $\alpha^n_{M_{H*}}$ determines an isomorphism of $H_1(M_H, \mathbb{Z})/_{torsion}$ and $H_1(M_{Ha}, \mathbb{Z})/_{torsion}$. Since also $\rho_a \circ r_* = \exp(\phi)$ with $\phi \in Hom(\pi_1(M_H), \mathbb{C})$ (see the definition of H), the above isomorphism implies that $\phi = (\eta + \phi_1) \circ \alpha^n_{M_{H*}}$, where ϕ_1 is a homomorphism of $\pi_1(M_{Ha})$ into $2\pi i\mathbb{Z}$. This shows that $\widetilde{\rho}_a$ is uniquely defined.

2. In fact the lemma is valid for any representation of $Hom(\pi_1(M), T_2(\mathbb{C}))$. However, our choice of the class is stipulated by the following argument.

Note that any ρ in Lemma 4.1 is equivalently defined by a flat vector bundle E on M,

$$0 \longrightarrow E_1 \longrightarrow E \longrightarrow E_2 \longrightarrow 0$$

where E_2 is a trivial flat vector bundle of complex rank-1 and E_1 is a flat vector bundle with unitary structure group (associated to the character ρ_a). It is well known that the equivalence classes of extensions of E_2 by E_1 are in one-to-one correspondence with the elements of the Čech cohomology group $H^1(M, \mathbf{E_1})$, where $\mathbf{E_1}$ is the sheaf of locally constant sections of E_1 (see, e.g., [A, Prop.2]). In particular, by the de Rham theorem and the Hodge decomposition we obtain that the equivalence class of E is uniquely defined by a d-harmonic 1-form η with values in E_1 (as before the Laplacian is defined by the flat Hermitian metric on E_1). The pullback r^*E on M_H determines the homomorphism $\rho|_{\pi_1(M_H)}$. Let $s : M_1 \longrightarrow M_{Ha}$ be a desingularization. (Since M_{Ha} is normal, the fibres of s are connected.) Then there is a modification $m : M'_H \longrightarrow M_H$ and a holomorphic surjective map $\alpha_1 : M'_H \longrightarrow M_1$ such that the diagram

$$\begin{tikzcd} M'_H \arrow[r, "\alpha_1"] \arrow[d, "m"'] & M_1 \arrow[d, "s"] \\ M_H \arrow[r, "\alpha^n_{M_{Ha}}"] & M_{Ha} \end{tikzcd} \tag{4.1}$$

commutes. In particular, Lemma 4.1 implies that there is a flat vector bundle F on M_1 lifted from M_{Ha} defined by

$$0 \longrightarrow F_1 \longrightarrow F \longrightarrow F_2 \longrightarrow 0$$

where F_2 is a trivial flat vector bundle of complex rank-1 and F_1 is a flat vector bundle with unitary structure group and by a d-harmonic 1-form η_1 with values in F_1 such that $\alpha_1^* F = m^*(r^*E)$ and, hence, $\alpha_1^*(\eta_1) = (r \circ m)^*(\eta)$. Note that according to part 1 of the remark, F_1 is uniquely defined by E_1. Clearly $\eta_1|_V = 0$ and $F_1|_V$ is trivial for any fibre V of s because $F|_V$ is a trivial flat vector bundle.

4.2. We are ready to prove Theorem 1.1. According to Proposition 2.3 it suffices to prove the theorem for representations $\rho \in Hom(\pi_1(M), T_n(\mathbb{C}))$. Let V_ρ be the associated flat vector bundle and $p_t : M_t \longrightarrow M$ be a regular covering with the covering group $T := Tor(\pi_1(M)/D\pi_1(M))$. The representation $\rho|_{\pi_1(M_t)}$ is defined by $p_t^* V_\rho$. Let $d(\rho) \in Hom(\pi_1(M), D_n(\mathbb{C}))$ be the representation defined by the diagonal of ρ. Then clearly $d(\rho)|_{\pi_1(M_t)} \in E_n(M_t)$ (see its definition in Introduction). Moreover, the diagonal characters ρ'_{ii} of $d(\rho)|_{\pi_1(M_t)}$ satisfy $\rho'_{ii} = \exp(\widetilde{\rho}_{ii} \circ p_{t*})$ for some $\widetilde{\rho}_{ii} \in Hom(\pi_1(M), \mathbb{C})$. In particular, $p_t^* V_\rho$ is defined by a flat connection ω on $M_t \times \mathbb{C}^n$ such that ω is an upper triangular 1-form and $diag(\omega)$ consists of harmonic 1-forms lifted from M. Further, according to the results of Section 2.2, $p_t^* V_\rho$ is also defined by a flat vector bundle E on M_t which is a direct sum of complex rank-1 flat vector bundles with structure group U_1 and by an upper triangular 1-form $\alpha + \partial h$ with values in $End(E)$ satisfying the flatness condition. Here α is d-harmonic with a nilpotent (0,1)-component and E is defined by the flat connection $diag(\omega'') - diag(\overline{\omega''})$ where ω'' is the (0,1)-component of ω. In particular, the structure of $diag(\omega)$ shows that $E = p_t^* V$ for a flat vector bundle V on M which is a direct sum $\oplus_{i=1}^n V_i$ of C^∞-trivial complex rank-1 flat vector bundles with structure group U_1. Further, $End(E) = \oplus_{i,j=1}^n E_{ij}$ with $E_{ij} := p_t^* V_i^* \otimes p_t^* V_j$. Then α admits a decomposition $\alpha = \sum_{i,j=1}^n \alpha_{ij}$ where α_{ij} is a harmonic 1-form with values in E_{ij}. Let $\mathcal{H}^1(E_{ij})$ be the space of E_{ij}-valued d-harmonic 1-forms on M_t. Since E_{ij} is pullback of a bundle on M, the finite abelian group T acts on $\mathcal{H}^1(E_{ij})$. This action is completely reducible and $\mathcal{H}^1(E_{ij})$ is isomorphic to the direct sum $\oplus_{k=1}^m H_k$ of unitary 1-dimensional T-modules. Then $\alpha_{ij} = \sum_{k=1}^m \alpha_{ij}^k$ with $\alpha_{ij}^k \in H_k$. Let $\xi_{ij}^k \in Hom(\pi_1(M_t), T_2^u)$ be a representation constructed by the pair (E_{ij}, α_{ij}^k).

Lemma 4.4 *There exists* $\widetilde{\xi}_{ij}^k \in Hom(\pi_1(M_{Ha}), T_2^u)$ *such that* $\xi_{ij}^k|_{\pi_1(M_H)} = (\alpha_{M_H}^n)^*(\widetilde{\xi}_{ij}^k)$.

Proof. According to our construction, for any $h \in T$ we have $h^*(\alpha_{ij}^k) = \rho_k(h)\alpha_{ij}^k$. Here we regard h as a biholomorphic map of M_t and ρ_k is the character determining H_k. In particular, ρ_k determines a complex rank-1 flat vector bundle B_k on M with structure group U_1 and α_{ij}^k determines a d-harmonic 1-form $\widetilde{\alpha}_{ij}^k$ with values in $B_k \otimes V_i^* \otimes V_j$ such that $p_t^*(\widetilde{\alpha}_{ij}^k) = \alpha_{ij}^k$. Let $\xi^k \in Hom(\pi_1(M), T_2)$ be a representation defined by $(B_k \otimes V_i^* \otimes V_j, \widetilde{\alpha}_{ij}^k)$. Then by Lemma 4.1 there is a representation $\widetilde{\xi}_{ij}^k \in Hom(\pi_1(M_{Ha}), T_2)$ such that $\xi^k \circ r_* = \widetilde{\xi}_{ij}^k \circ \alpha_{M_H *}^n$. But our constructions implies that $\xi^k \circ r_* = \xi_{ij}^k|_{\pi_1(M_H)}$.

The lemma is proved. $\square$

Further, observe that r^*E is the pullback of a bundle F defined on M_{Ha} lifted from $Alb(M)$ (because each V_i is defined by a homomorphism $\exp(\eta_i)$ with $\eta_i \in Hom(\pi_1(M), \mathbb{C})$). Moreover, F is the direct sum $\oplus_{i=1}^n F_i$ of C^∞-trivial complex rank-1 flat vector bundles with structure group U_1. Therefore $r^*(End(E))$ is the pullback of $End(F) = \oplus_{i,j=1}^n F_{ij}$, $F_{ij} := F_i^* \otimes F_j$ so that $r^* E_{ij} = (\alpha_{M_H}^n)^* F_{ij}$. Let $m : M'_H \longrightarrow M_H$ and s, α_1 be the same as in Remark 4.3 (see (4.1)). Then according to Lemma 4.4 and Remark 4.3 applied to each E_{ij} and α_{ij}^k, the form $(r \circ m)^*(\alpha_{ij})$ is the pullback by α_1 of a d-harmonic 1-form β_{ij} with values in $s^* F_{ij}$ (see also the uniqueness condition of part 1 of the remark) such that β_{ij} vanishes on each fibre of s. Thus $(r \circ m)^*(\alpha) = \alpha_1^*(\beta)$, where $\beta = \sum_{i,j=1}^n \beta_{ij}$ is a d-harmonic 1-form with values in $s^*(End(F))$. Observe also that the flatness condition and $\partial\overline{\partial}$-Lemma imply that $\alpha \wedge \alpha$ is a d-closed, d-exact (1,1)-form. Thus from Proposition 6.1 of [Br] it follows that $\beta \wedge \beta$ is a d-closed, d-exact matrix (1,1)-form with values in $End(F)$. Moreover, by the definition of α the (0,1)-component of β is nilpotent. Then from the results of Section 2.2 it follows that there is a unique (up to a flat additive summand) section h_1 such that $\beta + \partial h_1$ satisfies the flatness condition. Moreover, by the same uniqueness result applied to α and h we have

$$\alpha_1^*(\beta + \partial h_1) = (r \circ m)^*(\alpha + \partial h) \ .$$

In particular, the pair $(F, \beta + \partial h_1)$ determines a flat vector bundle F' on M_1 with triangular structure group such that $\alpha_1^* F' = (r \circ m)^* V_\rho$. Let V be a fibre of s (without loss of generality we may assume that V is smooth, for otherwise we apply the arguments below to each irreducible component of a desingularization of V). Then by the definition of β, $\beta|_V$ is zero and $F|_V$ is a trivial flat vector bundle. From here and the flatness condition it follows that $h_1|_V$ is pluriharmonic and so it is constant. Thus $F'|_V$ is a trivial flat vector bundle and the argument similar to that used in the proof of Proposition 3.3 shows that there is a flat vector bundle F'' defined on M_{Ha} such that $s^* F'' = F'$. Let $\widetilde{\rho} \in Hom(\pi_{M_{Ha}}, T_n(\mathbb{C}))$ be the homomorphism associated to F''. Then $\widetilde{\rho} \circ \alpha^n_{M_{H*}} = \rho \circ r_*$ because m_* determines an isomorphism of fundamental groups.

This completes the proof of Theorem 1.1. $\square$

Remark 4.5 Let $\rho \in Hom(\pi_1(M), N_n)$ be a homomorphism into the group of upper triangular unipotent matrices. Then the arguments of the proof of Theorem 1.1 show that $Ker(\alpha^n_{M*}) \subset Ker(\rho)$. In fact, in this case the above flat bundle E is the direct sum of trivial complex rank-1 flat vector bundles and so all harmonic 1-forms with values in $End(E)$ are pullbacks from $Alb(M)$. This result can be also obtained from a result of Campana (see e.g. [ABCKT, Prop.3.33]).

4.3. Proof of Theorem 1.4. Let $\xi \in S_n^{\mathcal{O}}(M)$. Condition (b) in the definition of the class $S_n^{\mathcal{O}}(M)$, the arguments used at the beginning of Section 4.2 and the results of Sections 2.2 imply that the equivalence class of $p_t^*(\xi)$ is uniquely defined by a d-harmonic matrix 1-form α with values in a bundle E which is a direct sum of C^∞-trivial complex rank-1 flat vector bundles with structure group U_1 lifted from M, such that $\alpha \wedge \alpha$ is d-exact (because the flat vector bundle on M_t associated to $p_t^*(\xi)$ is defined by a flat connection with an upper triangular (0,1)-component on $M_t \times \mathbb{C}^n$). Repeating word-for-word the arguments used in the proof of Theorem 1.1 in the case $\xi \in Hom(\pi_1(M), T_n(\mathbb{C}))$, we obtain the required factorization of ξ. It remains to prove that there is a $\rho \in Hom(\pi_1(M), D_n^u)$ such that $p_t^*(\xi) \in X_{p_t^*(\rho)}$. As above (see Section 2.2), $p_t^*(\xi)$ is defined by a matrix 1-form $(\alpha' + \partial h) + \alpha''$ with values in $End(E)$ satisfying the flatness condition. Here α' is a holomorphic 1-form and α'' is an antiholomorphic nilpotent 1-form. The flatness condition is equivalent to the equations

$$(\alpha' + \partial h) \wedge (\alpha' + \partial h) = 0;$$

$$\overline{\partial}(\alpha' + \partial h) = (\alpha' + \partial h) \wedge \alpha'' + \alpha'' \wedge (\alpha' + \partial h);$$

$$\alpha'' \wedge \alpha'' = 0 \ .$$

This implies that the form $z(\alpha' + \partial h) + \alpha''$ is also flat for any $z \in \mathbb{C}$. This form determines (by iterated path integration) a holomorphic deformation $\xi(z) \in Hom(\pi_1(M_t), GL_n(\mathbb{C}))$ such that $\xi(1) = p_t^*(\xi)$ and $\xi(0)$ is defined by $End(E)$-valued antiholomorphic nilpotent 1-form α''. In particular, $\xi(0) \in Hom(\pi_1(M_t), T_n^u)$ and the diagonal $d(\xi(0)) \in Hom(\pi_1(M_t), D_n^u)$ of $\xi(0)$ determines E. By the definition of E, there is a $\rho \in Hom(\pi_1(M), D_n^u)$ such that $p_t^*(\rho) = d(\xi(0))$. We complete the proof of the theorem by

Lemma 4.6 *Let Y be a compact Kähler manifold. For any $\eta \in Hom(\pi_1(Y), T_n(\mathbb{C}))$ we have $X_\eta \subset X_{d(\eta)}$.*

Proof. Let X_1 be an irreducible component of X_η of pure dimension ≥ 1. We will prove that $X_1 \subset X_{d(\eta)}$. According to the results of Section 2.3 there is an open neighbourhood $O \subset X_1$ of η such that for any $w \in O$, $w \neq \eta$, there is a holomorphic map $f : \mathbb{D} \longrightarrow X_1$ satisfying $f(0) = \eta$ and $w \in f(\mathbb{D})$. Regarding f as a holomorphic deformation of η we have $f(z)(g) = \sum_{i=0}^{\infty} f_i(g) z^i$, $g \in \pi_1(Y)$, such that $f(z) \in Hom(\pi_1(Y), GL_n(\mathbb{C}))$, $f_0 = \eta$, f_i are functions on $\pi_1(Y)$ with values in $gl_n(\mathbb{C})$ and the series converges uniformly on each compact of $\mathbb{D}$ for any $g \in \pi_1(Y)$. Let $y \in \mathbb{D}$, $y \neq 0$, be such that $f(y) = w$. For a holomorphic diagonal matrix function $A(z) := diag[1, z, z^2, ..., z^{n-1}]$ we define $f_1(z) = A(z)^{-1} f(y z^{n-1}) A(z)$. Set $a := (1/|y|)^{1/(n-1)} > 1$. Then a straightforward calculation shows that $f_1(z)$ is holomorphic in the disk $\mathbb{D}_a := \{z \in \mathbb{C} : |z| < a\}$, $f_1(0) = d(\eta)$, $f_1(1) = w$ and for any $z \in \mathbb{D}_a$, $f_1(z) \in Hom(\pi_1(Y), GL_n(\mathbb{C}))$. From here it follows that there is an irreducible component $X(w)$ of $X_{d(\eta)}$ containing w. This is valid for any $w \in O$. Thus X_1 coincides with $X(w_0)$ for some $w_0 \in O$ because X_1 is irreducible and $O \subset X_1$ is open.

This completes the proof of the lemma. □

The above lemma implies that $p_t^*(\xi)$ in Theorem 1.4 belongs to $X_{p_t^*(\rho)}$. □

Corollary 4.7 *Let $\rho \in S_n(M)$ (see the definition in Introduction). Then there is a holomorphic deformation $\rho(z) \in Hom(\pi_1(M), T_n(\mathbb{C}))$, $z \in \mathbb{C}$, of ρ such that $\rho(1) = \rho$ and $\rho(0) \in Hom(\pi_1(M), D_n^u)$.*

Proof. The proof repeats the arguments of the proof of Theorem 1.4, because in this case ρ is defined by an upper triangular flat connection ω on $M \times \mathbb{C}^n$ and we can use the results of Section 2.2. Further details might be left to the reader. □

Proof of Theorem 1.6. Clearly we have $G_s^0(M) \subseteq G_{su}^0(M)$. We will prove that $G_{su}^0(M) \subseteq G_s^0(M)$. Let $\rho \in S_n(M)$, that is $\rho \in Hom(\pi_1(M), T_n(\mathbb{C}))$ and $d(\rho) \in E_n$. According to Corollary 4.7, there is a holomorphic deformation $\rho(z) \in$

$Hom(\pi_1(M), T_n(\mathbb{C}))$, $z \in \mathbb{C}$, of ρ such that $\rho(1) = \rho$, $\rho(0) \in Hom(\pi_1(M), D_n^u)$ and $\rho(z)(g) = \sum_{i=0}^{\infty} f_i(g) z^i$, $g \in \pi_1(M)$ where the f_i are upper triangular functions on $\pi_1(M)$. Let $\pi_m : GL_n(\mathbb{C})[[z]] \longrightarrow GL_n(A_m[z])$ be the quotient homomorphism defined in Section 2.3. Then $\pi_m(\rho(z)) \in Hom(\pi_1(M), GL_n(A_m[z]))$ is equivalent to a representation $\widetilde{\rho} \in Hom(\pi_1(M), T_p^u)$ (see the arguments of the proof of Proposition 2.3). Therefore $Ker(\pi_m(\rho(z))) \supset G_{su}^0(M)$ for any $m \geq 0$. Then Proposition 2.2 implies that

$$G_{su}^0(M) \subset \bigcap_{i \geq 0} Ker(\pi_i(\rho(z))) = \bigcap_{z \in \mathbb{C}} Ker(\rho(z)) \subset Ker(\rho) \ .$$

This shows that $G_{su}^0(M) \subset G_s^0(M)$ and completes the proof of the theorem. □

5. Proofs of Theorem 1.7 and Other Results.

According to Theorem 1.1, $Ker(\alpha_{M_{H*}}^n) \subset G_s(M)$. We assume that the Albanese map α_{M_H} is non-trivial. For otherwise, $Ker(\alpha_{M_{H*}}^n) = \pi_1(M_H)$ which implies that $M_s = M_H$ and the required statement is trivial. Let $K := Ker(\alpha_{M_{H*}})$. Then K is a normal subgroup of $\pi_1(M)$ (because $\pi_1(M_H) \subset \pi_1(M)$ is normal) and $\pi_1(M)/K$ is a solvable group of the form

$$\{e\} \longrightarrow \mathbb{Z}^{2l} \longrightarrow \pi_1(M)/K \longrightarrow H \longrightarrow \{e\} \ .$$

Here $l = dim_{\mathbb{C}} Alb(M_H)$. It is easy to see that $\pi_1(M)/K$ admits a faithful representation to some $T_n(\mathbb{C})$ (see, e.g. the construction used in Lemma 4.4). Therefore $G_s(M) \subset K$. Let $F := \alpha_{M_{H*}}^n(G_s(M))$ and $M_{Fa} \longrightarrow M_{Ha}$ be a covering corresponding to the group $F \subset \pi_1(M_{Ha})$. Here $M_{Ha} = \alpha_{M_H}^n(M_H)$. By the covering homotopy theorem there is a holomorphic surjective map $f : M_s \longrightarrow M_{Fa}$ which covers $\alpha_{M_H}^n$. Let $i : V \hookrightarrow M_H$ be a connected component of a fibre of $\alpha_{M_H}^n$. Then $i_*(\pi_1(V)) \subset Ker(\alpha_{M_{H*}}^n) \subset G_s(M)$ and therefore there is an embedding $i' : F \hookrightarrow M_s$ such that $i'(F)$ is a connected component of a fibre of f. This shows that f is a proper surjective holomorphic map onto M_{Fa}. Let $f = f_1 \circ f_2$ be the Stein factorization of f. Here $f_2 : M_s \longrightarrow M'$ is a proper holomorphic map with connected fibres and $f_1 : M' \longrightarrow M_{Fa}$ is a finite morphism. Then M' is normal and any holomorphic function on M_s is pullback of a holomorphic function on M'. We will prove that M' is holomorphically convex.

Let $K' = \alpha_{M_{H*}}^n(K)$ and $M_{K'a}$ be the Galois covering of M_{Ha} corresponding to K'. Then by the definition of K, there is an intermediate covering map $p : M_{Fa} \longrightarrow M_{K'a}$. Let $A : \mathbb{C}^l \longrightarrow Alb(M_H)$ be the universal covering. Since by definition $\pi_1(M_H)/K \cong \pi_1(M_{Ha})/K' \cong \pi_1(Alb(M_H))$, $M_{K'a}$ is normalization of $A^{-1}(\alpha_{M_H}(M_H)) \subset \mathbb{C}^l$. In particular, the standard properties of Stein spaces (see e.g. [GR]) imply that $M_{K'a}$ is Stein and the covering M_{Fa} of $M_{K'a}$ is also a normal Stein space. Moreover, $f_1 : M' \longrightarrow M_{Fa}$ is a finite morphism and so M' is a normal Stein space. In particular, M' is holomorphically convex.

Other statements of the theorem are easy corollaries of the above argument.

The theorem is proved. □

Remark 5.1 According to Remark 1.2 the kernel of any matrix unipotent representation of $\pi_1(M)$ contains $Ker(\alpha^n_{M*})$. Then the same arguments as before show that the Malcev covering of M is holomorphically convex which includes the result of Katzarkov [Ka1].

Proof of Corollary 1.8. If $G_s(M) = \{e\}$ then the universal covering M_u is holomorphically convex and any fibre V of $\alpha^n_{M_H}$ can be lifted to M_u (we denote the image of V in M_u by the same letter). Since $\pi_2(M) = 0$, Hurewicz's theorem implies that $H^2(M_u, \mathbb{C}) = 0$. Let ω be a d-closed (1,1)-form on M_u which is pullback of the form on M determining the Kähler metric. If $dim_{\mathbb{C}} V \geq 1$ then $\omega|_V$ is not d-exact and so ω determines a non-trivial element of $H^2(M_u, \mathbb{C})$. This contradiction shows that $dim_{\mathbb{C}} V = 0$ and the map $f_2 : M_u \longrightarrow M'$ from the proof of Theorem 1.7 is identity. Therefore $M_u = M'$ is Stein. $\square$

Remark 5.2 In particular, M_u satisfying assumptions of Corollary 1.8 has the structure of a k-dimensional complex with $k = \dim_{\mathbb{C}} M_u$. Thus any compact complex manifold M satisfying the assumptions of the corollary such that $\pi_i(M) \neq 0$ for some $i > dim_{\mathbb{C}} M$ is not Kähler.

Proof of Theorem 1.12. According to Corollary 1.8, $\alpha^n_{M_H} : M_H \longrightarrow M_{Ha}$ is a finite analytic covering. In particular,

$$dim_C M_H \leq dim_{\mathbb{C}} Alb(M_H) \leq \frac{1}{2} rank(\pi_1(M)/D^2\pi_1(M)) \; .$$

These inequalities and the condition of the theorem imply that

$$dim_C M_H = dim_{\mathbb{C}} Alb(M_H) = \frac{1}{2} rank(\pi_1(M)/D^2\pi_1(M))$$

and $\alpha_{M_H} : M_H \longrightarrow Alb(M_H)$ is a finite analytic covering. According to Theorem 1.4, any $\rho \in Hom(\pi_1(M), T_n(\mathbb{C}))$ factors through $\alpha^n_{M_H}$ (which coincides with α_{M_H} in our case). Therefore $\pi_1(M_H)/G_s(M) \cong \pi_1(Alb(M_H))$. Since $G_s(M) = \{e\}$, we have $\pi_1(M_H) \cong \pi_1(Alb(M_H)) \cong \mathbb{Z}^{2l}$. This completes the proof of the first part of the theorem.

If, in addition, $\pi_i(M_u) = 0$ for $1 \leq i \leq dim_{\mathbb{C}} M$, then M_H is a $K(\mathbb{Z}^{2l}, 1)$-space. Note that $Alb(M_H)$ is also a $K(\mathbb{Z}^{2l}, 1)$-space and α_{M_H} induces an isomorphism of fundamental groups. Therefore by Eilenberg's theorem (see [E]), $\alpha^*_{M_H} : H^i(Alb(M_H), \mathbb{Z}) \longrightarrow H^i(M_H, \mathbb{Z})$ is an isomorphism for any $i \geq 0$. Note that if $1 \in H^{2l}(Alb(M_H), \mathbb{Z})$ is a generator then $\alpha^*_{M_H}(1)$ coincides with $deg(\alpha_{M_H})$. Applying the above isomorphisms with $i = 2l$ we get $deg(\alpha_{M_H}) = 1$ showing that α_{M_H} is biholomorphic. $\square$

Similarly one can prove the statement of Remark 1.13.

References

[ABCKT] J.Amorós, M.Burger, K.Corlette, D.Kotschick, and D.Toledo, Fundamental groups of compact Kähler manifolds, Math. Surveys Monographs, **44**, Amer. Math. Soc., Providence, RI, 1996.

[A] M.F.Atiyah, Complex analytic connections in fibre bundles, Trans. Amer. Math. Soc. **85** (1957), 181-207.

[Be] A.Beauville, Annulation du H^1 pour les fibrés en droites plats, LNM, **1507**, Springer Verlag (1992), 1-15.

[BK] F.Bogomolov and L.Katzarkov, Complex projective surfaces and infinite groups, GAFA, vol. 8 (1998), 243-272.

[BO] A.Brudnyi and A.Onishchik, Triangular de Rham cohomology of compact Kähler manifolds, To appear in Math. Sbornik, **2**, 2001.

[BS] A.I.Borevich and I.R.Shafarevich, Number theory, Translated from Russian. Pure and Applied Mathematics, vol. 20, Academic Press, New York, 1966.

[Br] A.Brudnyi, Classification theorem for a class of flat connections and representations of Kähler groups, Michigan Math. J., **46** (3) (1999), 489-514.

[C] F.Campana, Remarques sur le groupes de Kähleriennes nilpotent, Ann. Scien. L'Ecole Norm. Sup. **28**, fasc. 3 (1995), 307-316.

[C1] F.Campana, Ensembles de Green-Lazarsfeld et quotients resolubles des groupes de Kähler, preprint, 2000.

[E] S.Eilenberg, Homology of spaces with operators 1, Trans.Amer. Math.Soc., **61** (1947), 378-417.

[G] A. Guichardet, Cohomologie des groupes topologiques et des algèbres de Lie, Cedic/Fernand Nathan, Paris (1980).

[GR] H.Grauert and R.Remmert, Theorie der Steinschen Räume, Springer-Verlag (1977).

[Ka1] L.Katzarkov, Nilpotent groups and universal coverings of smooth projective varieties, J. Differential Geometry, **45** (1997), 336-348.

[Ka2] L.Katzarkov, On the Shafarevich maps, Proc. Symp. in Pure Math., Proceedings of AMS 62:2 (1997), 173-217.

[KaR] L.Katzarkov and M.Ramachandran, On the universal coverings of algebraic surfaces, Annales Scien. E.N.S. **31** (1998), 887-888.

[KN] S.Kobayashi and K.Nomizu, Foundations of differential geometry, vol.1, Interscience, New York, 1963.

[Ko1] J.Kollár, Shafarevich maps and plurigenera of algebraic varieties, Inventiones Matematicae **113**, Fasc. 1, no.1 (1993), 177-215.

[Ko2] J.Kollár, Shafarevich maps and automorphic forms, Princeton University Press, Princeton, NJ (1995).

[L] S. Lojasiewicz, Triangulation of semi-analytic sets, Ann. Scuola Norm. Sup. Pisa, (3) **18** (1964), 449–474.

[M] D. Mumford, Abelian varieties, Lect. in Tata Inst. of Fundamental Research, Bombay (1968).

[N] T.Napier, Convexity properties of coverings of smooth projective varieties, Mathematische Annalen, **286** (1990), 433-479.

[O] A.L.Onishchik, Certain concepts and applications of non-Abelian cohomology theory, Trudy Moskov. Mat. Obshch. **17** (1967), 45-88.

[S] C.T.Simpson, Constructing variations of Hodge structures using Yang-Mills theory and applications to unifarmization, J. Amer. Math. Soc., 1:4 (1998), 867-918.

[S1] C.T.Simpson, Subspaces of Moduli Spaces of rank one local systems, Ann. Sc. ENS, **26** (1993), 361-401.

[Y] S.T.Yau, Uniformization of geometric structures, Proc. of Symp. in Pure Mathematics, **48**, AMS (1988), 265-274.